



# Baryon spectroscopy and structure in the Dyson-Schwinger approach

**Gernot Eichmann**

IST Lisboa, Portugal

**WE Heraeus seminar “From correlation functions to QCD phenomenology”  
Bad Honnef, Germany**

**April 3, 2018**

# Why?

**QCD Lagrangian:**  $\mathcal{L} = \bar{\psi} (\not{\partial} + ig\mathbf{A} + m) \psi + \frac{1}{4} F_{\mu\nu}^a F_a^{\mu\nu}$

- if it only were that simple...  
we don't measure quarks and gluons, but **hadrons**



mesons



baryons



glueballs?



hybrids?



tetraquarks?



pentaquarks??

- origin of **mass generation** and **confinement?**

	u	d	s	c	b	t
Current mass [GeV]	0.003	0.005	0.1	1	4	175
„Constituent“ mass [GeV]	0.35	0.35	0.5	1.5	4.5	175

- need to understand **spectrum and interactions!**

# The hadron zoo

## Mesons

$0^-$	$0^{++}$	$1^-$	$1^{--}$	$1^{++}$	$1^{+-}$	$2^-$	$2^{++}$	$3^-$
$\pi(140)$ $\pi(1300)$ $\pi(1800)$	$a_0(980)$ $a_0(1450)$ $a_0(1960)$	$\pi_1(1400)$ $\pi_1(1600)$	$\rho(770)$ $\rho(1450)$ $\rho(1570)$ $\rho(1700)$ $\rho(1900)$	$a_1(1260)$ $a_1(1420)$ $a_1(1640)$	$h_1(1235)$	$\pi_2(1670)$ $\pi_2(1880)$	$a_2(1320)$ $a_2(1700)$	$\rho_2(1690)$ $\rho_2(1990)$
$K(494)$ $K(1460)$ $K(1830)$	$K_0^*(800)$ $K_0^*(1430)$ $K_0^*(1960)$	$K^*(892)$ $K^*(1410)$ $K^*(1680)$	$K_1(1400)$ $K_1(1650)$	$K_1(1270)$ $K_2(1680)$ $K_2(1770)$ $K_2(1820)$	$K_0^*(1430)$ $K_0^*(1980)$	$K_3(1780)$	$\omega(782)$ $\phi(1020)$ $\omega(1420)$ $\omega(1680)$ $\phi(1680)$	$f_1(1285)$ $f_1(1420)$ $f_1(1810)$
$\eta(548)$ $\eta'(958)$ $\eta(1296)$ $\eta(1405)$ $\eta(1475)$ $\eta(1760)$	$f_0(500)$ $f_0(980)$ $f_0(1370)$ $f_0(1500)$ $f_0(1710)$	$\omega(1270)$ $h_1(1170)$ $h_1(1380)$ $h_1(1595)$	$\eta_2(1645)$ $\eta_2(1870)$	$f_2(1270)$ $f_2(1430)$ $f_2'(1825)$ $f_2(1680)$ $f_2(1640)$ $f_2(1810)$ $f_2(1910)$ $f_2(1960)$	$\omega_2(1670)$ $\phi_2(1850)$			




## Baryons


$\frac{1}{2}^+$	$\frac{1}{2}^-$	$\frac{3}{2}^+$	$\frac{3}{2}^-$	$\frac{5}{2}^+$	$\frac{5}{2}^-$	$\frac{7}{2}^+$
$N(939)$ $N(1440)$ $N(1710)$ $N(1880)$	$N(1535)$ $N(1650)$ $N(1895)$	$N(1720)$ $N(1900)$	$N(1520)$ $N(1700)$ $N(1875)$	$N(1680)$ $N(1860)$ $N(2000)$	$N(1675)$	$N(1900)$
$\Delta(1910)$	$\Delta(1620)$ $\Delta(1900)$	$\Delta(1232)$ $\Delta(1800)$ $\Delta(1920)$	$\Delta(1700)$ $\Delta(1840)$ $\Delta(2000)$	$\Delta(1905)$ $\Delta(2000)$	$\Delta(1980)$	$\Delta(1950)$
$\Lambda(1116)$ $\Lambda(1800)$ $\Lambda(1810)$	$\Lambda(1405)$ $\Lambda(1670)$ $\Lambda(1800)$	$\Lambda(1890)$	$\Lambda(1520)$ $\Lambda(1680)$	$\Lambda(1820)$	$\Lambda(1830)$	
$\Sigma(1189)$ $\Sigma(1660)$ $\Sigma(1880)$	$\Sigma(1760)$	$\Sigma(1385)$	$\Sigma(1670)$ $\Sigma(1940)$	$\Sigma(1915)$	$\Sigma(1775)$	
$\Xi(1315)$	$\Xi(1530)$	$\Xi(1820)$	$\Xi(1672)$			



# Light baryons



- Extraction of resonances?



- Gluon exchange** vs. flavor dependence?
- Nature of **Roper**?
- qqq vs. **quark-diquark**?


- “Quark core” vs. **chiral dynamics**?
- Admixture of **multiquarks**?
- Hybrid baryons**?

# Hadrons in QCD

**Lattice:** extract baryon poles from (gauge-invariant) two-point correlators:

$$G(x-y) = \langle 0 | T \underbrace{[\Gamma_{\alpha\beta\gamma} \psi_{\alpha} \psi_{\beta} \psi_{\gamma}]}_{B(x)}(x) \underbrace{[\bar{\Gamma}_{\rho\sigma\tau} \bar{\psi}_{\rho} \bar{\psi}_{\sigma} \bar{\psi}_{\tau}]}_{\bar{B}(y)}(y) | 0 \rangle = \int \mathcal{D}[\psi, \bar{\psi}, A] e^{-S} B(x) \bar{B}(y)$$

$$G(\tau) \sim e^{-m\tau} \quad \Leftrightarrow \quad G(P^2) \sim \frac{1}{P^2 + m^2}$$




- **Infinite volume:**  
Bound states, resonances,  
branch cuts


- **Finite volume:**  
bound states & scattering states

# Hadrons in QCD


**Lattice:** extract baryon poles from (gauge-invariant) two-point correlators:

$$G(x-y) = \langle 0 | T \underbrace{[\Gamma_{\alpha\beta\gamma} \psi_{\alpha} \psi_{\beta} \psi_{\gamma}]}_{B(x)}(x) \underbrace{[\bar{\Gamma}_{\rho\sigma\tau} \bar{\psi}_{\rho} \bar{\psi}_{\sigma} \bar{\psi}_{\tau}]}_{\bar{B}(y)}(y) | 0 \rangle = \int \mathcal{D}[\psi, \bar{\psi}, A] e^{-S} B(x) \bar{B}(y)$$

$$= \lim_{\substack{x_i \rightarrow x \\ y_i \rightarrow y}} \Gamma_{\alpha\beta\gamma} \bar{\Gamma}_{\rho\sigma\tau} \langle 0 | T \psi_{\alpha}(x_1) \psi_{\beta}(x_2) \psi_{\gamma}(x_3) \bar{\psi}_{\rho}(y_1) \bar{\psi}_{\sigma}(y_2) \bar{\psi}_{\tau}(y_3) | 0 \rangle$$


$$= x \otimes \text{---} G \text{---} \otimes y \xrightarrow{P^2 \rightarrow -m_A^2} \text{---} \otimes \text{---} \text{---} \otimes y$$


Alternative: extract **gauge-invariant** baryon poles from **gauge-fixed** quark 6-point function:



**Bethe-Salpeter wave function:**


residue at pole, contains all information about baryon

$$\langle 0 | T \psi_{\alpha}(x_1) \psi_{\beta}(x_2) \psi_{\gamma}(x_3) | \lambda \rangle$$

# QCD's n-point functions


QCD's classical action:

$$S = \int d^4x \left[ \bar{\psi} (\not{\partial} + ig\mathcal{A} + m) \psi + \frac{1}{4} F_{\mu\nu}^a F_a^{\mu\nu} \right]$$




Quantum "effective action":

$$\int \mathcal{D}[\psi, \bar{\psi}, A] e^{-S} = e^{-\Gamma}$$



DSEs = quantum equations of motion:

derived from path integral, relate n-point functions



- infinitely many coupled equations
- reproduce perturbation theory, but **nonperturbative**
- systematic truncations: neglect higher n-point functions to obtain **closed system**

Some Reviews:

Roberts, Williams, *Prog. Part. Nucl. Phys.* 33 (1994),


Alkofer, von Smekal, *Phys. Rept.* 353 (2001)

GE, Sanchis-Alepuz, Williams, Alkofer, Fischer,

*Prog. Part. Nucl. Phys.* 91 (2016), 1606.09602 [hep-ph]


# QCD's n-point functions

## • Quark propagator



**Dynamical chiral symmetry breaking** generates 'constituent-quark masses'

## • Gluon propagator





## • Three-gluon vertex

$F_1 [ \delta^{\mu\nu} (p_1 - p_2)^\rho + \delta^{\nu\rho} (p_2 - p_3)^\mu + \delta^{\rho\mu} (p_3 - p_1)^\nu ] + \dots$

Agreement between lattice, DSE & FRG within reach

→ looking forward to the talks at this workshop!

## • Quark-gluon vertex







# DSEs $\rightarrow$ Hadrons?

## Bethe-Salpeter approach:


use scattering equation  $G = G_0 + G_0 K G$



- still exact - to begin with, kernel is black box
- but can be derived together with QCD's n-point functions. Important to preserve symmetries!




Homogeneous BSE for **BS wave function**:




# DSEs $\rightarrow$ Hadrons?

## Bethe-Salpeter approach:


use scattering equation  $G = G_0 + G_0 K G$




- still exact - to begin with, kernel is black box
- but can be derived together with QCD's n-point functions. Important to preserve symmetries!



Homogeneous BSE for **BS wave function**




... or **BS amplitude**:




# Mesons

- Meson **Bethe-Salpeter equation** in QCD:




$$K(M) G(M) \phi_i(M) = \lambda_i(M) \phi_i(M)$$


- Depends on QCD's n-point functions, satisfy **DSEs**:




- Kernel derived in accordance with **chiral symmetry**:



- Eigenvalues in **pion** channel:



- Quark propagator has **complex singularities**: no physical threshold




# Mesons


- Meson **Bethe-Salpeter equation** in QCD:

$$K(M) G(M) \phi_i(M) = \lambda_i(M) \phi_i(M)$$

- Depends on QCD's n-point functions, satisfy **DSEs**:



- Kernel derived in accordance with **chiral symmetry**:




**Rainbow-ladder:**  
effective gluon exchange

$$\alpha(k^2) = \alpha_{\text{IR}}\left(\frac{k^2}{\Lambda^2}, \eta\right) + \alpha_{\text{UV}}(k^2)$$


adjust scale  $\Lambda$  to observable,  
keep width  $\eta$  as parameter

Maris, Tandy, PRC 60 (1999),  
Qin et al., PRC 84 (2011)

Eigenvalues in **pion channel**:




Quark propagator has **complex singularities**: no physical threshold




# Mesons

- Pion is **Goldstone boson**:  $m_\pi^2 \sim m_q$



- Light meson spectrum** beyond rainbow-ladder




Williams, Fischer, Heupel, PRD 93 (2016)

GE, Sanchis-Alepuz, Williams, Alkofer, Fischer, PNP 91 (2016)


see also Chang, Roberts, PRL 103 (2009), PRC 85 (2012)

- Charmonium spectrum**

Fischer, Kubrak, Williams, EPJ A 51 (2015)



- Pion transition form factor**




GE, Fischer, Weil, Williams, PLB 774 (2017)

# Baryons

## Covariant Faddeev equation for baryons:


GE, Alkofer, Krassnigg, Nicmorus, PRL 104 (2010)



- 3-gluon diagram vanishes  $\Rightarrow$  **3-body effects small?**

Sanchis-Alepuz, Williams, PLB 749 (2015)


- 2-body kernels same as for mesons, no further approximations:



$$\Psi_{\alpha\beta\gamma\delta}(p, q, P) = \sum_i f_i(p^2, q^2, p \cdot q, p \cdot P, q \cdot P) \tau_i(p, q, P)_{\alpha\beta\gamma\delta}$$


Lorentz-invariant  
dressing functions

Dirac-Lorentz  
tensors carry  
OAM: s, p, d,...



Review: GE, Sanchis-Alepuz, Williams, Alkofer, Fischer,  
PPNP 91 (2016), 1606.09602

# Form factors




$$J^\mu = e \bar{u}(p_f) \left( F_1(Q^2) \gamma^\mu + F_2(Q^2) \frac{i}{4m} [\gamma^\mu, \not{Q}] \right) u(p_i)$$

## Consistent derivation of **current matrix elements & scattering amplitudes**

Kvinikhidze, Blankleider, PRC 60 (1999),  
GE, Fischer, PRD 85 (2012) & PRD 87 (2013)

$$J^\mu = \text{[diagram 1]} + \text{[diagram 2]} + \text{[diagram 3]} + \text{[diagram 4]}$$



- **rainbow-ladder** topologies (1st line):



- **quark-photon vertex** preserves em. gauge invariance, dynamically generates **VM poles**:


$$\text{[diagram 1]} = \text{[diagram 2]} \rightarrow \text{[diagram 3]}$$

# Form factors

## Nucleon em. form factors from three-quark equation

GE, PRD 84 (2011)





- “Quark core without pion cloud”



- **similar:**  $N \rightarrow \Delta\gamma$  transition, axial & pseudoscalar FFs, octet & decuplet em. FFs

Review: GE, Sanchis-Alepuz, Williams, Fischer, Alkofer, PPNP 91 (2016), 1606.09602

$$J^\mu = \text{[Diagram 1]} + \text{[Diagram 2]} + \text{[Diagram 3]} + \text{[Diagram 4]}$$





# Scattering amplitudes


## Scattering amplitudes from quark level:

### • $\pi\pi$ scattering

DSE: Bicudo, Cotanch, Llanes-Estrada, Maris, Ribeiro, Szczepaniak, PRD 65 (2002),

Cotanch, Maris, PRD 66 (2002)

CST: Biernat, Pena, Ribeiro, Stadler, Gross, PRD 90 (2014)



Universal band

◆ ChPT tree, 1 loop, 2 loops

■ ChPT + dispersion theory (2001)

□ DIRAC (2005)

■ NA48  $K \rightarrow 3\pi$  (2005)

■ E865 isospin corrected

■ NA48 isospin-corrected

■ MILC (2004)


■ NPLQCD (2005)

■ Del Debbio (2007)

■ ETM (2007)

★ DSE (rainbow-ladder)


### • Nucleon Compton scattering



GE, Fischer, PRD 85 (2012) & PRD 87 (2013), GE, FBS 57 (2016)



### • Hadronic light-by-light scattering

Goecke, Fischer, Williams, PLB 704 (2011),  
GE, Fischer, Heupel, PRD 92 (2015)



# The role of diquarks


Mesons and 'diquarks' closely related:  
 after taking traces, only factor 1/2 remains  
 ⇒ **diquarks 'less bound' than mesons**



**Pseudoscalar & vector mesons**  
 already good in rainbow-ladder

**Scalar & axialvector mesons**  
 too light, repulsion beyond RL

↔





**Scalar & axialvector diquarks**  
 sufficient for nucleon and  $\Delta$

↔

**Pseudoscalar & vector diquarks**  
 important for remaining channels

# Baryon spectrum


Quark-diquark with reduced pseudoscalar + vector diquarks: [GE, Fischer, Sanchis-Alepuz, PRD 94 \(2016\)](#)




- Scale  $\Lambda$  set by  $f_\pi$
- Current-quark mass  $m_q$  set by  $m_\pi$
- $c$  adjusted to  $\rho$ - $a_1$  splitting
- $\eta$  doesn't change much

# Strange baryons


	[nn]	{nn}	[ns]	{ns}	{ss}
$N$	●	●			
$\Delta$		●			
$\Lambda$	●		●	●	
$\Sigma$		●	●	●	
$\Xi$			●	●	●
$\Omega$					●



# Strange baryons







**N**




**$\Omega$**


# Strange baryons




# Strange baryons



**N**



**Δ**



**Λ**

- Strange baryons similar to **light baryons**:

$$\begin{aligned} \Omega &\rightarrow \Delta \\ \Sigma, \Xi &\rightarrow N + \Delta \quad \rightarrow \text{rich spectrum!} \\ \Lambda &\rightarrow N + \text{singlets} \end{aligned}$$

- Roper,  $\Delta(1600)$ ,  $\Lambda(1405)$ ,  $\Lambda(1520)$ : additional dynamics?

GE, Fischer, in preparation


# The role of diquarks?

- **Singlet:** symmetric variable, carries overall scale:

$$S_0 \sim p_1^2 + p_2^2 + p_3^2 + \frac{M^2}{3}$$


- **Doublet:**  $\mathcal{D}_0 \sim \frac{1}{S_0} \begin{bmatrix} -\sqrt{3}(\delta x + 2\delta\omega) \\ x + 2\omega \end{bmatrix}$

Mandelstam plane,  
outside: **diquark poles!**



Lorentz invariants can be grouped into **multiplets of the permutation group S3:**

GE, Fischer, Heupel, PRD 92 (2015)



- **Second doublet:**  $\mathcal{D}_1 \sim \frac{1}{\sqrt{S_0}} \begin{bmatrix} -\sqrt{3}(\delta x - \delta\omega) \\ x - \omega \end{bmatrix}$

$f_i(S_0, \text{red circle}, \text{red circle}) \rightarrow$  full result as before



$f_i(S_0, \text{pink circle}, \text{pink circle}) \rightarrow$  **same ground-state spectrum,**  
but diquark poles switched off!




# Scattering amplitudes

## TPE corrections to form factors

Guichon, Vanderhaeghen, PRL 91 (2003)




- **Nucleon Compton scattering**



GE, Fischer, PRD 85 (2012) &  
PRD 87 (2013), GE, FBS 57 (2016)

## Proton radius puzzle?

Antonigni et al., 2013, Pohl et al. 2013,  
Birse, McGovern 2012, Carlson 2015




## Nucleon polarizabilities

Hagelstein, Miskimen, Pascalutsa,  
Prog. Part. Nucl. Phys. 88 (2016)

## Structure functions & PDFs in forward limit

$$\text{Diagram} = \sum \text{Diagram} \sim \left| \text{Diagram} \right|^2$$

## Handbag dominance & GPDs in DVCS




## $p\bar{p}$ annihilation

@ PANDA

# Compton scattering


Compton amplitude = sum of **Born terms** + 1PI structure part:



Griesshammer, McGovern, Phillips, Feldman, Prog. Part. Nucl. Phys. 67 (2012)


# Compton scattering

Scattering amplitude: [GE, Fischer, PRD 85 \(2012\) & PRD 87 \(2013\)](#)



- **Poincaré covariance** and **crossing symmetry** automatic
- **em. gauge invariance** and **chiral symmetry** automatic  
as long as all ingredients calculated from symmetry-preserving kernel
- **perturbative processes** included
- **s, t, u channel poles** dynamically generated,  
no need for “offshell hadrons”

# Nucleon resonances



$J^P = \frac{1}{2}^+$	$\frac{3}{2}^+$	$\frac{1}{2}^-$	$\frac{3}{2}^-$
N(940) N(1440) N(1710) N(1880)	N(1720) N(1900)	N(1535) N(1650) N(1895)	N(1520) N(1700) N(1875)
Δ(1910)	Δ(1232) Δ(1600) Δ(1920)	Δ(1620) Δ(1900)	Δ(1700) Δ(1940)

## Need em. transition FFs

But vertices are half offshell:  
need 'consistent couplings'

[Pascalutsa, Timmermans, PRC 60 \(1999\)](#)

- **em gauge invariance:**  $Q^\mu \Gamma^{\alpha\mu} = 0$
- **spin-3/2 gauge invariance:**  $k^\alpha \Gamma^{\alpha\mu} = 0$
- invariance under **point transformations:**  $\gamma^\alpha \Gamma^{\alpha\mu} = 0$
- no kinematic dependencies, **"minimal" basis**


E.g. Jones-Scadron current  
**cannot be used offshell:**

$$\Gamma^{\alpha\mu} \sim \bar{u}^\alpha(k) \left[ m^2 \lambda_- (G_M^* - G_E^*) \varepsilon_{kQ}^{\alpha\mu} - G_E^* \varepsilon_{kQ}^{\alpha\beta} \varepsilon_{kQ}^{\beta\mu} - \frac{1}{2} G_C^* Q^\alpha k^\beta t_{QQ}^{\beta\mu} \right] u(k')$$

$$t_{AB}^{\alpha\beta} = A \cdot B \delta^{\alpha\beta} - B^\alpha A^\beta$$

$$\varepsilon_{AB}^{\alpha\beta} = \gamma_5 \varepsilon^{\alpha\beta\gamma\delta} A^\gamma B^\delta$$

# Nucleon resonances



$J^P = \frac{1}{2}^+$	$\frac{3}{2}^+$	$\frac{1}{2}^-$	$\frac{3}{2}^-$
N(940) N(1440) N(1710) N(1880)	N(1720) N(1900)	N(1535) N(1650) N(1895)	N(1520) N(1700) N(1875)
Δ(1910)	Δ(1232) Δ(1600) Δ(1920)	Δ(1620) Δ(1900)	Δ(1700) Δ(1940)

## Need em. transition FFs

But vertices are half offshell:  
need 'consistent couplings'

Pascalutsa, Timmermans, PRC 60 (1999)

- **em gauge invariance:**  $Q^\mu \Gamma^{\alpha\mu} = 0$
- **spin-3/2 gauge invariance:**  $k^\alpha \Gamma^{\alpha\mu} = 0$
- invariance under **point transformations:**  $\gamma^\alpha \Gamma^{\alpha\mu} = 0$
- no kinematic dependencies, **"minimal" basis**

## Most general offshell vertices


satisfying these constraints:

GE, Ramalho, in preparation

$$\frac{1}{2}^+ \rightarrow \frac{1}{2}^\pm : \Gamma^\mu = \begin{bmatrix} 1 \\ \gamma_5 \end{bmatrix} \sum_{i=1}^8 F_i T_i^\mu \left\{ \begin{array}{l} t_{QQ}^{\mu\nu} \gamma^\nu \\ [\gamma^\mu, \not{Q}] \\ \dots \end{array} \right.$$

$$\frac{1}{2}^+ \rightarrow \frac{3}{2}^\pm : \Gamma^{\alpha\mu} = \begin{bmatrix} \gamma_5 \\ 1 \end{bmatrix} \sum_{i=1}^{12} F_i T_i^{\alpha\mu} \left\{ \begin{array}{l} \epsilon_{kQ}^{\alpha\mu} \\ t_{kQ}^{\alpha\mu} \\ it_{k\gamma}^{\alpha\beta} t_{QQ}^{\beta\mu} \\ \dots \end{array} \right.$$

# Nucleon resonances




$J^P = \frac{1}{2}^+$	$\frac{3}{2}^+$	$\frac{1}{2}^-$	$\frac{3}{2}^-$
N(940) N(1440) N(1710) N(1880)	N(1720) N(1900)	N(1535) N(1650) N(1895)	N(1520) N(1700) N(1875)
Δ(1910)	Δ(1232) Δ(1600) Δ(1920)	Δ(1620) Δ(1900)	Δ(1700) Δ(1940)

## Need em. transition FFs


But vertices are half offshell:  
need 'consistent couplings'

[Pascalutsa, Timmermans, PRC 60 \(1999\)](#)



- **em gauge invariance:**  $Q^\mu \Gamma^{\alpha\mu} = 0$
- **spin-3/2 gauge invariance:**  $k^\alpha \Gamma^{\alpha\mu} = 0$
- invariance under **point transformations:**  $\gamma^\alpha \Gamma^{\alpha\mu} = 0$
- no kinematic dependencies, **"minimal" basis**



# Nucleon resonances





$J^P = \frac{1}{2}^+$	$\frac{3}{2}^+$	$\frac{1}{2}^-$	$\frac{3}{2}^-$
N(940) N(1440) N(1710) N(1880)	N(1720) N(1900)	N(1535) N(1650) N(1895)	N(1520) N(1700) N(1875)
Δ(1910)	Δ(1232) Δ(1600) Δ(1920)	Δ(1620) Δ(1900)	Δ(1700) Δ(1940)





Example:  
N(1535) helicity amplitudes

- PDG
- CLAS data  
[userweb.jlab.org/~mokeev/resonance\\_electrocouplings](http://userweb.jlab.org/~mokeev/resonance_electrocouplings)
- MAID  
Tiator, Drechsel, Kamalov, Vanderhaeghen, EPJ 198 (2011)

# Nucleon resonances



**N(1535) transition FFs:**  
no kinematic constraints





Example:  
**N(1535) helicity amplitudes**

- PDG
- CLAS data  
[userweb.jlab.org/~mokeev/resonance\\_electrocouplings](http://userweb.jlab.org/~mokeev/resonance_electrocouplings)
- MAID  
Tiator, Drechsel, Kamalov, Vanderhaeghen, EPJ 198 (2011)





# Nucleon resonances



**N(1535) transition FFs:**  
no kinematic constraints

Fit



Example:  
**N(1535) helicity amplitudes**

PDG


CLAS data

[userweb.jlab.org/~mokeev/resonance\\_electrocouplings](http://userweb.jlab.org/~mokeev/resonance_electrocouplings)

MAID

Tiator, Drechsel, Kamalov, Vanderhaeghen, EPJ 198 (2011)

# Nucleon resonances



$$J^P = \frac{1}{2}^+$$

$$\frac{3}{2}^+$$

$$\frac{1}{2}^-$$

$$\frac{3}{2}^-$$

**N(940)**

**N(1720)**

**N(1535)**

**N(1520)**

**N(1440)**

*N(1900)*

**N(1650)**

*N(1700)*

*N(1710)*

*N(1895)*

*N(1875)*

*N(1880)*

**$\Delta(1910)$**

**$\Delta(1232)$**

**$\Delta(1620)$**

**$\Delta(1700)$**

$\Delta(1600)$

$\Delta(1900)$

$\Delta(1940)$

$\Delta(1920)$

# Kinematics



$$= \sum_{i=1}^{18} c_i(\eta_+, \eta_-, \omega, \lambda) \bar{u}(p_f) X_i^{\mu\nu}(p, Q, Q') u(p_i)$$

## 18 CFFs

4 kinematic variables:

$$\eta_+ = \frac{Q^2 + Q'^2}{2m^2}$$

$$\eta_- = \frac{Q \cdot Q'}{m^2}$$

$$\omega = \frac{Q^2 - Q'^2}{2m^2}$$

$$\lambda = -\frac{p \cdot Q}{m^2}$$

## 18 Compton tensors, form minimal basis

- systematic derivation
- similar to Tarrach basis

[Tarrach, Nuovo Cim. A28 \(1975\)](#)

$$X'_i = U_{ij} X_j, \quad \det U = \text{const.}$$

- CFFs free of kinematics

$$X_1^{\mu\nu} = \frac{1}{m^4} t_{Q'p}^{\mu\alpha} t_{pQ}^{\alpha\nu},$$

$$X_2^{\mu\nu} = \frac{1}{m^2} t_{Q'Q}^{\mu\nu},$$

$$X_3^{\mu\nu} = \frac{1}{m^4} t_{Q'Q'}^{\mu\alpha} t_{QQ}^{\alpha\nu},$$

$$X_4^{\mu\nu} = \frac{1}{m^6} t_{Q'Q'}^{\mu\alpha} p^\alpha p^\beta t_{QQ}^{\beta\nu},$$





$$X_5^{\mu\nu} = \frac{\lambda}{m^4} (t_{Q'Q'}^{\mu\alpha} t_{pQ}^{\alpha\nu} + t_{Q'p}^{\mu\alpha} t_{QQ}^{\alpha\nu}),$$

$$X_6^{\mu\nu} = \frac{1}{m^2} \varepsilon_{Q'Q}^{\mu\nu},$$

$$X_7^{\mu\nu} = \frac{1}{im^3} (t_{Q'Q'}^{\mu\alpha} \varepsilon_{\gamma Q}^{\alpha\nu} - \varepsilon_{Q'\gamma}^{\mu\alpha} t_{QQ}^{\alpha\nu}),$$




$$X_8^{\mu\nu} = \frac{\omega}{im^3} (t_{Q'Q'}^{\mu\alpha} \varepsilon_{\gamma Q}^{\alpha\nu} + \varepsilon_{Q'\gamma}^{\mu\alpha} t_{QQ}^{\alpha\nu}),$$

⋮




GE, Ramalho,  
in preparation

# Compton form factors




To be multiplied with 
$$\frac{(m_R^2 - m^2)^2}{(s - m_R^2)(u - m_R^2)} = \frac{\delta^2}{(\eta_- + \delta)^2 - 4\lambda^2}$$

# Compton form factors




- CS on scalar particle

# Compton form factors




- CS on scalar particle
- CS on pointlike scalar

# Compton form factors




- CS on scalar particle
- CS on pointlike scalar
- CS on pointlike fermion

# Compton form factors






# Compton form factors




- CS on scalar particle
- CS on pointlike scalar
- CS on pointlike fermion
- Nucleon Born poles in s & u channel
- **Scalar pole** in t channel

# Compton form factors





GE, Fischer, Weil, Williams,  
PLB 774 (2017)

- CS on scalar particle
- CS on pointlike scalar
- CS on pointlike fermion
- Nucleon Born poles in s & u channel
- Scalar pole in t channel
- Pion pole in t channel ( $\pi^0 \rightarrow \gamma^* \gamma^*$ )





# Polarizabilities



Scalar polarizabilities:

$$\begin{bmatrix} \alpha + \beta \\ \beta \end{bmatrix} = -\frac{\alpha_{\text{em}}}{m^3} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$$




Spin polarizabilities:

$$\begin{bmatrix} \gamma_{E1E1} \\ \gamma_{M1M1} \\ \gamma_{E1M2} \\ \gamma_{M1E2} \end{bmatrix} = \frac{\alpha_{\text{em}}}{2m^4} \begin{bmatrix} c_6 + 4c_{11} - 4c_{12} \\ -c_6 - 2c_{10} + 4c_{12} \\ c_6 + 2c_{10} \\ -c_6 \end{bmatrix}$$


$$\begin{bmatrix} \gamma_0 \\ \gamma_\pi \end{bmatrix} = -\frac{2\alpha_{\text{em}}}{m^4} \begin{bmatrix} c_{11} \\ c_6 + c_{10} + c_{11} - 2c_{12} \end{bmatrix}$$

# Polarizabilities



Scalar polarizabilities:

$$\begin{bmatrix} \alpha + \beta \\ \beta \end{bmatrix} = -\frac{\alpha_{\text{em}}}{m^3} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$$




Hagelstein, Miskimen, Pascalutsa,  
Prog. Part. Nucl. Phys. 88 (2016)

PDG:


$$-c_1 = 20.3(4)$$

$$-c_2 = 3.7(6)$$

Large  $\Delta(1232)$  contribution,  
but also  $N(1520)$  non-negligible



# Spin polarizabilities




Only  $\Delta(1232)$   
important

# Compton scattering




- kinematic variables
- tensor basis
- constraint-free **Compton FFs**


GE, Ramalho,  
in preparation



- general offshell transition vertices
- constraint-free **transition FFs**
- fits for transition FFs




- impact of higher resonances on Compton FFs
- only  $\Delta(1232)$  and  $N(1520)$  relevant for polarizabilities



# Developing numerical tools




→ poster by [Esther Weil](#)

Rare pion decay  $\pi^0 \rightarrow e^+e^-$ :



$$A(t) = \int d\sigma \int dz \dots \frac{1}{k^2+m^2} \frac{1}{Q^2} \frac{1}{Q'^2}$$

Photon and lepton poles produce branch cuts in complex plane:  
**deform integration contour!**




- Result agrees with dispersion relations
- Algorithm is stable & efficient
- Can be applied to any integral as long as **singularity locations** known  
[Weil, GE, Fischer, Williams, PRD 96 \(2017\)](#)
- Integrate behind quark singularities!  
[Windisch, PRC 95 \(2017\)](#)

# Tetraquarks

→ poster by **Paul Wallbott**

- **Light scalar mesons  $\sigma$ ,  $\kappa$ ,  $a_0$ ,  $f_0$  as tetraquarks:**  
solution of four-body equation reproduces mass pattern

GE, Fischer, Heupel, PLB 753 (2016)




BSE dynamically generates  
**meson poles** in wave function:

$$f_i(S_0, \nabla, \triangle, \circ) \rightarrow 1500 \text{ MeV}$$


$$f_i(S_0, \nabla, \triangle, \circ) \rightarrow 1500 \text{ MeV}$$

$$f_i(S_0, \nabla, \triangle, \circ) \rightarrow 1200 \text{ MeV}$$

$$f_i(S_0, \nabla, \triangle, \circ) \rightarrow \mathbf{350 \text{ MeV !!}}$$




Four quarks rearrange  
to "**meson molecule**"



- Similar in **meson-meson / diquark-antidiquark** approximation  
(analogue of quark-diquark for baryons)


Heupel, GE, Fischer, PLB 718 (2012)






# Towards multiquarks

Transition from **quark-gluon** to **nuclear degrees of freedom**:




- 6 ground states, one of them **deuteron**  
[Dyson, Xuong, PRL 13 \(1964\)](#)
- Dibaryons vs. **hidden color**?  
[Bashkanov, Brodsky, Clement, PLB 727 \(2013\)](#)
- **Deuteron FFs** from quark level?

**Microscopic origins of nuclear binding?**



[Weise, Nucl. Phys. A805 \(2008\)](#)




- only quarks and gluons
- **quark interchange** and **pion exchange** automatically included
- **dibaryon** exchanges

---


# Backup slides

# Resonances?

$\rho \rightarrow \pi\pi$ : **resonance dynamics**  
 only beyond rainbow-ladder,  
 would shift  $\rho$  pole into complex plane  
 (above  $\pi\pi$  threshold)





But  $\rho$  decay width  
 already calculable  
 in rainbow-ladder



## Rainbow-ladder vs. lattice:

References: GE et al., PNP 91 (2016) 1606.09602




actual resonance dynamics  
 subleading effect?


$\rho$  may just be a special case,  
 but baryon spectrum?

# Resonances?

$\rho \rightarrow \pi\pi$ : **resonance dynamics**  
 only beyond rainbow-ladder,  
 would shift  $\rho$  pole into complex plane  
 (above  $\pi\pi$  threshold)





But  $\rho$  decay width  
 already calculable  
 in rainbow-ladder



## Rainbow-ladder vs. lattice:


References: GE et al., PNP 91 (2016) 1606.09602




actual resonance dynamics  
 subleading effect?

$\rho$  may just be a special case,  
 but baryon spectrum?

# Nucleon resonances



# Nucleon resonances



# Nucleon resonances







$\Delta(1232)$

CLAS

PDG

MAID

Fit



# Nucleon resonances







$\Delta(1700)$

CLAS

PDG

MAID

Fit





# Nucleon resonances







N(1520)

CLAS

PDG

MAID

Fit




# Bethe-Salpeter equations

## Simplest: Wick-Cutkosky model


Wick 1954, Cutkosky 1954, Nakanishi 1969, ...

- scalar tree-level propagators, scalar exchange particle
- bound states for  $M < 2m$


$$K(M) G(M) \phi_i(M) = \lambda_i(M) \phi_i(M)$$

But:

- no confinement: threshold  $2m$
- not a consistent QFT: would need to solve DSEs for propagators, vertices etc.




# Bethe-Salpeter equations

## Simplest: Wick-Cutkosky model



Wick 1954, Cutkosky 1954, Nakanishi 1969, ...

- scalar tree-level propagators, scalar exchange particle
- bound states for  $M < 2m$



But:

- no confinement: threshold  $2m$
- not a consistent QFT: would need to solve DSEs for propagators, vertices etc.



# Bethe-Salpeter equations

## Simplest: Wick-Cutkosky model


Wick 1954, Cutkosky 1954, Nakanishi 1969, ...

- scalar tree-level propagators, scalar exchange particle
- bound states for  $M < 2m$

$$K(M) G(M) \phi_i(M) = \lambda_i(M) \phi_i(M)$$

But:


- no confinement: threshold  $2m$
- not a consistent QFT: would need to solve DSEs for propagators, vertices etc.



# Form factors



## Nucleon charge radii:

isovector (p-n) Dirac (F1) radius




## Nucleon magnetic moments:

isovector (p-n), isoscalar (p+n)



- **Pion-cloud effects** missing ( $\Rightarrow$  divergence!), agreement with lattice at larger quark masses.



- **But:** pion-cloud cancels in  $\kappa^S \Leftrightarrow$  quark core

Exp:  $\kappa^S = -0.12$

Calc:  $\kappa^S = -0.12(1)$



GE, PRD 84 (2011)


# Lattice vs. DSE / BSE

## Lattice

### Full dynamics

contained in path integral

Proper treatment of  
**resonances** essential



Simpler access to **position-space**  
and **gluonic operators**

$$\langle N | \bar{\psi} \not{D} \psi | N \rangle \sim \text{diagram 1} + \text{diagram 2}$$


Two Feynman diagrams representing the nucleon axial current. Each diagram shows a blue oval representing a nucleon with two external lines. A wavy line representing a gluon is attached to the top vertex of the nucleon. The first diagram shows the gluon attached to the left vertex, and the second diagram shows it attached to the right vertex.

**Precision!**

## DSE / BSE

Dynamics constructed from  
underlying **n-point functions**

Resonance dynamics  
“on top of” **quark-gluon dynamics**



Simpler access to multi-scale problems  
and higher n-point functions



Can tell us about underlying dynamics!

# nPI effective action

nPI effective actions provide **symmetry-preserving closed truncations**.

3PI at 3-loop: **all two- and three-point functions are dressed**; 4, 5, ... do not appear.

$$\Gamma_2 = - \text{[circle with dashed line and dot]} + \frac{1}{2} \text{[circle with dashed line and two dots]} + \frac{1}{4} \text{[circle with dashed line and four dots]}$$

see: Sanchis-Alepuz & Williams, J. Phys. Conf. Ser. 631 (2015), arXiv:1503.05896 and refs therein

**Self-energy:**

$$\Sigma = \frac{\delta\Gamma_2}{\delta D} = - \text{[dashed arc]} - \text{[dashed arc]} + \text{[dashed arc]} + \text{[dashed arc]} = - \text{[dashed arc]}$$

**Vertex:**

$$\frac{\delta\Gamma_2}{\delta V} = 0 \Rightarrow - \text{[vertex]} + \text{[vertex]} + \text{[vertex]} = 0$$

**Vacuum polarization:**

$$\Sigma' = \frac{\delta\Gamma_2}{\delta D'} = - \text{[circle with dashed line]} + \frac{1}{2} \text{[circle with dashed line]} + \frac{1}{2} \text{[circle with dashed line]} = - \frac{1}{2} \text{[circle with dashed line]}$$

**BSE kernel:**

$$-K = \frac{\delta\Sigma}{\delta D} = - \text{[kernel]} - \text{[kernel]} + \text{[kernel]} + \text{[kernel]} + \text{[kernel]} + \text{[kernel]} = - \text{[kernel]} + \text{[kernel]}$$

# nPI effective action

nPI effective actions provide **symmetry-preserving closed truncations**.

3PI at 3-loop: **all two- and three-point functions are dressed**; 4, 5, ... do not appear.

$$\Gamma_2 = - \text{[circle with dashed line]} + \frac{1}{2} \text{[circle with dashed line]} + \frac{1}{4} \text{[circle with dashed line and two internal vertices]}$$

see: Sanchis-Alepuz & Williams,  
J. Phys. Conf. Ser. 631 (2015), arXiv:1503.05896 and refs therein

So we arrive at a closed system of equations:

- Crossed ladder cannot be added by hand, requires **vertex correction!**



# nPI effective action

nPI effective actions provide **symmetry-preserving closed truncations**.

3PI at 3-loop: **all two- and three-point functions are dressed**; 4, 5, ... do not appear.

$$\Gamma_2 = - \text{[circle with dashed line]} + \frac{1}{2} \text{[circle with dashed line and dot]} + \frac{1}{4} \text{[crossed ladder]}$$

see: Sanchis-Alepuz & Williams,  
J. Phys. Conf. Ser. 631 (2015), arXiv:1503.05896 and refs therein

So we arrive at a closed system of equations:


$$\begin{aligned} \text{[solid line with dot]}^{-1} &= \text{[solid line]}^{-1} + \text{[crossed ladder]} \\ \text{[dashed line with dot]}^{-1} &= \text{[dashed line]}^{-1} + \frac{1}{2} \text{[circle with dot]} \\ \text{[3-point vertex]} &= \text{[tree-level vertex]} - \text{[rainbow-ladder]} \\ \text{[2-point function]} &= \text{[rainbow-ladder]} - \text{[crossed ladder]} \end{aligned}$$

- Crossed ladder cannot be added by hand, requires **vertex correction**!
- without 3-loop term: **rainbow-ladder** with tree-level vertex  $\Rightarrow$  2PI
- but still requires **DSE solutions** for propagators!
- Similar in QCD. nPI truncation guarantees chiral symmetry, massless pion in chiral limit, etc.

# Baryon spectrum I

Three-quark vs. quark-diquark in rainbow-ladder: [GE, Fischer, Sanchis-Alepuz, PRD 94 \(2016\)](#)

M [GeV]





- **qqg** and **q-dq** agrees: N, Δ, Roper, N(1535)
- # levels compatible with experiment: **no states missing**
- N, Δ and their 1st excitations (including **Roper**) agree with experiment
- But remaining states too low  $\Rightarrow$  wrong level ordering between Roper and N(1535)

# Baryon spectrum

Quark-diquark with reduced pseudoscalar + vector diquarks: [GE, Fischer, Sanchis-Alepuz, PRD 94 \(2016\)](#)


M [GeV]



- Scale  $\Lambda$  set by  $f_\pi$
- Current-quark mass  $m_q$  set by  $m_\pi$
- $c$  adjusted to  $\rho$ - $a_1$  splitting
- $\eta$  doesn't change much



# Baryon spectrum

M [GeV]



Level ordering between  
**Roper and N(1535):**


dynamics of ps diquark produces  
2 nearby states: **N(1535), N(1650)**



# Resonances

- **Current-mass evolution of Roper:**


GE, Fischer, Sanchis-Alepuz, PRD 94 (2016)




- **'Pion cloud' effects difficult to implement at quark-gluon level:**



- Branch cuts & widths generated by **meson-baryon interactions**: Roper  $\rightarrow N\pi$ , etc.



- **Lattice: finite volume, DSE (so far): bound states**



Resonance dynamics shifts poles into complex plane, but effects on real parts small?

# QED


## QED's classical action:

$$S = \int d^4x \left[ \bar{\psi} (\not{\partial} + ig\mathbf{A} + m) \psi + \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \right]$$



## Quantum "effective action":


$$\int \mathcal{D}[\psi, \bar{\psi}, A] e^{-S} = e^{-\Gamma}$$



**Perturbation theory:** expand Green functions in powers of the coupling


$$\frac{1}{A(p^2)(i\not{p} + M(p^2))} = \frac{1}{i\not{p} + m} + \text{loop diagrams} + \dots$$

mass function



$$\frac{1}{D^{-1}(p^2)(p^2 \delta^{\mu\nu} - p^\mu p^\nu)} = \frac{1}{p^2 \delta^{\mu\nu} - p^\mu p^\nu} + \text{loop diagrams} + \dots$$

running coupling



$$F_1 \gamma^\mu - \frac{F_2}{2m} \sigma^{\mu\nu} Q^\nu + \dots = \gamma^\mu + \text{loop diagrams} + \dots$$

anomalous magnetic moment  
 $F_2(0) = \frac{\alpha}{2\pi}$

# QED


## QED's classical action:

$$S = \int d^4x \left[ \bar{\psi} (\not{\partial} + ig\mathbf{A} + m) \psi + \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \right]$$




## Quantum "effective action":

$$\int \mathcal{D}[\psi, \bar{\psi}, A] e^{-S} = e^{-\Gamma}$$



**Perturbation theory:** expand Green functions in powers of the coupling



$\Rightarrow$  extremely precise theory predictions!





# Dynamical quark mass

Simplest example: **Munczek-Nemirovsky model**  
Gluon propagator =  $\delta$ -function, analytically solvable

Munczek, Nemirovsky, PRD 28 (1983)

$$D^{\mu\nu}(k) \Gamma^\nu(p, q) \longrightarrow \sim \Lambda^2 \delta^4(k) \gamma^\mu$$

Quark DSE becomes

$$S^{-1}(p) - S_0^{-1}(p) = \Lambda^2 \gamma^\mu S(p) \gamma^\mu = \Lambda^2 \frac{2i\not{p} + 4M}{(p^2 + M^2)A},$$

leads to self-consistent equations for **A**, **M**:

$$A = 1 + \frac{2\Lambda^2}{(p^2 + M^2)A}, \quad AM = m_0 + 2M \frac{2\Lambda^2}{(p^2 + M^2)A}$$

Two solutions in chiral limit: IR + UV


$$\begin{aligned} M(p^2) &= \sqrt{\Lambda^2 - p^2} & M(p^2) &= 0 \\ A(p^2) &= 2 & A(p^2) &= \frac{1}{2} \left( 1 + \sqrt{1 + 8\Lambda^2/p^2} \right) \end{aligned}$$

Quark condensate:

$$-\langle \bar{q}q \rangle = N_C \int \frac{d^4p}{(2\pi)^4} \text{Tr} S(p) = \frac{2}{15} \frac{N_C}{(2\pi)^2} \Lambda^3$$

$$S(p) = \frac{1}{A(p^2)} \frac{-i\not{p} + M(p^2)}{p^2 + M^2(p^2)}$$

$$S^{-1}(p) = A(p^2) (i\not{p} + M(p^2))$$




Another extreme case: **NJL model**,  
gluon propagator = const,  
 $M(p^2) = \text{const}$ , but critical behavior

Nambu, Jona-Lasinio, 1961


# Dynamical quark mass

- Simplest realistic example: **rainbow-ladder**



Tree-level quark-gluon vertex + **effective interaction**:

$$D^{\mu\nu}(k) \Gamma^\nu(p, q) \rightarrow \sim \frac{\alpha(k^2)}{k^2} \left( \delta^{\mu\nu} - \frac{k^\mu k^\nu}{k^2} \right) \gamma^\nu$$




$$\alpha(k^2) = \alpha_{\text{IR}} \left( \frac{k^2}{\Lambda^2}, \eta \right) + \alpha_{\text{UV}}(k^2)$$

adjust scale  $\Lambda$  to observable,  
keep width  $\eta$  as parameter

Maris, Tandy, PRC 60 (1999)

- If strength is large enough ( $\alpha > \alpha_{\text{crit}}$ ): **DCSB**
- All dimensionful quantities  $\sim \Lambda$  in chiral limit  
 $\Rightarrow$  **mass generation for hadrons!**



Classical PCAC relation for  $SU(N_f)_A$ :

$$\partial_\mu \bar{\psi} \gamma^\mu \gamma_5 \mathbf{t}_a \psi = i \bar{\psi} \{ M, \mathbf{t}_a \} \gamma_5 \psi$$


At quantum level:

$$f_\pi m_\pi^2 = 2m r_\pi$$


Also  $f_\pi \sim \Lambda \Rightarrow m_\pi = 0$  in chiral limit!  
 $\Rightarrow$  **massless Goldstone bosons!**

# Extracting resonances

## Hadronic coupled-channel equations:




Sato-Lee/EBAC/ANL-Osaka, Dubna-Mainz-Taiwan, Valencia, Jülich-Bonn, GSI, JLab, MAID, SAID, KSU, Giessen, Bonn-Gatchina, JPAC, ...



Suzuki et al., PRL 104 (2010)


## Microscopic effects?

What is an “offshell hadron”?



# Extracting resonances


Photoproduction of **exotic mesons** at JLab/GlueX:



What if exotic mesons are **relativistic  $q\bar{q}$  states**?  
 $\Rightarrow$  study with DSE/BSE!


**Scattering amplitudes** at quark-gluon level:

GE, Fischer, PRD 85 (2012), PRD 87 (2013)




# Diquarks?

- Suggested to resolve ‘**missing resonances**’ in quark model:  
fewer degrees of freedom  $\Rightarrow$  fewer excitations



Anselmino et al., *Rev. Mod. Phys.* 65 (1993),  
Klempt, Richard, *Rev. Mod. Phys.* 82 (2010)

- QCD version: assume  $qq$  scattering matrix as sum of diquark correlations  
 $\Rightarrow$  three-body equation simplifies to **quark-diquark BSE**



Oettel, Alkofer, Hellstern Reinhardt, *PRC* 58 (1998),  
Cloet, GE, El-Bennich, Klähn, Roberts, *FBS* 46 (2009)

**Quark exchange** binds nucleon, gluons absorbed in building blocks.  
Scalar diquark  $\sim 800$  MeV, axialvector diquark  $\sim 1$  GeV

Maris, *FBS* 32 (2002), GE, Krassnigg, Schwinzerl, Alkofer, *Ann. Phys.* 323 (2008), GE, *FBS* 57 (2016)

- N and  $\Delta$  properties similar in quark-diquark and three-quark approach:  
**quark-diquark approximation is good!**

# Complex eigenvalues?




**Excited states:** some EVs are complex conjugate?

Typical for **unequal-mass** systems, already in Wick-Cutkosky model

Wick 1954, Cutkosky 1954

Connection with “**anomalous**” states?

Ahlig, Alkofer, Ann. Phys. 275 (1999)



If  $G = G^\dagger$  and  $G > 0$  :  
Cholesky decomposition  $G = L^\dagger L$

$$K L^\dagger L \phi_i = \lambda_i \phi_i$$

$$(L K L^\dagger) (L \phi_i) = \lambda_i (L \phi_i)$$

$\Rightarrow$  Hermitian problem with same EVs!

$K$  and  $G$  are Hermitian (even for unequal masses!) but  $KG$  is not

# Complex eigenvalues?



**Excited states:** some EVs are complex conjugate?

Typical for **unequal-mass** systems, already in Wick-Cutkosky model

Wick 1954, Cutkosky 1954

Connection with “**anomalous**” states?

Ahlig, Alkofer, *Ann. Phys.* 275 (1999)



$K$  and  $G$  are Hermitian (even for unequal masses!) but  $KG$  is not

If  $G = G^\dagger$  and  $G > 0$  :  
Cholesky decomposition  $G = L^\dagger L$

$$K L^\dagger L \phi_i = \lambda_i \phi_i \quad \Rightarrow \text{Hermitian problem with same EVs!}$$

$$(LKL^\dagger)(L\phi_i) = \lambda_i (L\phi_i)$$

- $\Rightarrow$  all EVs strictly **real**
- $\Rightarrow$  level repulsion
- $\Rightarrow$  “anomalous states” removed?

# Complex eigenvalues?


**Excited states:** some EVs are complex conjugate?

Typical for **unequal-mass** systems, already in Wick-Cutkosky model


Wick 1954, Cutkosky 1954

Connection with **“anomalous” states**?

Ahlig, Alkofer, *Ann. Phys.* 275 (1999)



$K$  and  $G$  are Hermitian (even for unequal masses!) but  $KG$  is not



Eigenvalue spectrum for pion channel

GE, FBS 58 (2017)

— only **pos.** EVs in  $G$   
 - - - only **neg.** EVs in  $G$

If  $G = G^\dagger$  and  $G > 0$ :

Cholesky decomposition  $G = L^\dagger L$

$$K L^\dagger L \phi_i = \lambda_i \phi_i$$

$$(L K L^\dagger) (L \phi_i) = \lambda_i (L \phi_i)$$

$\Rightarrow$  Hermitian problem with same EVs!

$\Rightarrow$  all EVs strictly **real**



$\Rightarrow$  level repulsion

$\Rightarrow$  “anomalous states” removed?



# Tetraquarks?

Light scalar ( $0^{++}$ ) mesons don't fit into the conventional meson spectrum:



$f_0$  (980 MeV)  $s\bar{s}$   
 $\kappa$  (680 MeV)  $u\bar{s}, d\bar{s}$   
 $a_0$  (980 MeV) }  $u\bar{u}, d\bar{d}, u\bar{d}$   
 $\sigma$  (500 MeV)

- Why are  $a_0, f_0$  mass-degenerate?
- Why are their **decay widths** so different?



$$\Gamma(\sigma, \kappa) \approx 550 \text{ MeV}$$

$$\Gamma(a_0, f_0) \approx 50\text{--}100 \text{ MeV}$$

- Why are they so **light**?  
 Scalar mesons  $\sim$  **p-waves**, should have masses similar to axialvector & tensor mesons  $\sim 1.3 \text{ GeV}$

# Tetraquarks?

What if they were **tetraquarks** (diquark-antidiquark)? [Jaffe 1977](#), [Close, Tornqvist 2002](#), [Maiani, Polosa, Riquer 2004](#)



$f_0$  (980 MeV) }  $us\bar{u}s, \dots$   
 $a_0$  (980 MeV) }  
 $\kappa$  (800 MeV) }  $us\bar{u}d, \dots$   
 $\sigma$  (500 MeV) }  $ud\bar{u}d$


- Explains **mass ordering & decay widths**:  
 $f_0$  and  $a_0$  couple to  $K\bar{K}$ , large widths for  $\sigma, \kappa$

- Alternative: **meson molecules?**  
[Weinstein, Isgur 1982, 1990](#); [Close, Isgur, Kumano 1993](#)

- Non- $q\bar{q}$  nature** of  $\sigma$  supported by dispersive analyses, unitarized ChPT, large  $N_c$ , extended linear  $\sigma$  model, quark models  
[Pelaez, Phys. Rept. 658 \(2016\)](#)



# Four-body equation



**Two-body interactions**

... plus permutations:

$$(qq)(\bar{q}\bar{q}), (q\bar{q})(q\bar{q}), (q\bar{q})(q\bar{q})$$

$$(12)(34) \quad (23)(14) \quad (13)(24)$$

**3-body**  
(+ permutations)

**4-body**

**Bethe-Salpeter amplitude:**

$$\Gamma(p, q, k, P) = \sum_i f_i(p^2, q^2, k^2, \{\omega_j\}, \{\eta_j\}) \tau_i(p, q, k, P) \otimes \text{Color} \otimes \text{Flavor}$$

**9 Lorentz invariants:**

$$p^2, \quad q^2, \quad k^2$$

$$\omega_1 = p \cdot k \quad \eta_1 = p \cdot P$$

$$\omega_2 = p \cdot k \quad \eta_2 = q \cdot P$$

$$\omega_3 = p \cdot q \quad \eta_3 = k \cdot P$$

$$P^2 = -M^2$$

**256**  
**Dirac-**  
**Lorentz**  
**tensors**

**2 Color**  
**tensors:**

$$3 \otimes \bar{3}, \quad 6 \otimes \bar{6} \quad \text{or}$$

$$1 \otimes 1, \quad 8 \otimes 8$$

(Fierz-equivalent)


# Structure of the amplitude

- **Singlet:** symmetric variable, carries overall scale:

$$S_0 = \frac{1}{4} (p^2 + q^2 + k^2)$$

- **Doublet:**  $\mathcal{D}_0 = \frac{1}{4S_0} \begin{bmatrix} \sqrt{3}(q^2 - p^2) \\ p^2 + q^2 - 2k^2 \end{bmatrix}$

Mandelstam triangle,  
outside: **meson and diquark poles!**




Lorentz invariants can be grouped into **multiplets of the permutation group S4**:

GE, Fischer, Heupel, PRD 92 (2015)

- **Triplet:**  $\mathcal{T}_0 = \frac{1}{4S_0} \begin{bmatrix} 2(\omega_1 + \omega_2 + \omega_3) \\ \sqrt{2}(\omega_1 + \omega_2 - 2\omega_3) \\ \sqrt{6}(\omega_2 - \omega_1) \end{bmatrix}$

tetrahedron bounded by  $p_i^2 = 0$ ,  
outside: **quark singularities**



- **Second triplet:**  
3dim. sphere

$$\mathcal{T}_1 = \frac{1}{4S_0} \begin{bmatrix} 2(\eta_1 + \eta_2 + \eta_3) \\ \sqrt{2}(\eta_1 + \eta_2 - 2\eta_3) \\ \sqrt{6}(\eta_2 - \eta_1) \end{bmatrix}$$

# Tetraquark mass

$$f_i(\mathcal{S}_0, \nabla, \triangle, \circ)$$

Idea: use symmetries to figure out **relevant** momentum dependence


similar:

- **Three-gluon vertex**

[GE, Williams, Alkofer, Vujanovic, PRD 89 \(2014\)](#)


- **HLbL scattering for muon g-2**

[GE, Fischer, Heupel, PRD 92 \(2015\)](#)




# Tetraquark mass

$$f_i(\mathcal{S}_0, \nabla, \triangle, \circ)$$




# Tetraquark mass

$$f_i(\mathcal{S}_0, \nabla, \triangle, \circ)$$



# Tetraquark mass



$$f_i(\mathcal{S}_0, \nabla, \triangle, \circ)$$






# Tetraquark mass

$$f_i(S_0, \nabla, \triangle, \circ)$$




# Towards multiquarks

Transition from **quark-gluon** to **nuclear degrees of freedom**:




- 6 ground states, one of them **deuteron**  
[Dyson, Xuong, PRL 13 \(1964\)](#)
- Dibaryons vs. **hidden color**?  
[Bashkanov, Brodsky, Clement, PLB 727 \(2013\)](#)
- **Deuteron FFs** from quark level?

**Microscopic origins of nuclear binding?**




[Weise, Nucl. Phys. A805 \(2008\)](#)



- only quarks and gluons
- **quark interchange** and **pion exchange** automatically included
- **dibaryon** exchanges


# Hadron physics with functional methods

Understand properties of  
**elementary n-point functions**



↔



Calculate hadronic **observables**:  
mass spectra, form factors, scattering amplitudes, . . .



- **QCD**
- **symmetries** intact (Poincare invariance & chiral symmetry important)
- access to all momentum scales & all quark masses
- compute mesons, baryons, tetraquarks, . . . **from same dynamics**
  
- **systematic** construction of truncations
- technical challenges: coupled integral equations, complex analysis, structure of 3-, 4-, ... point functions, **need lots of computational power!**


**access to underlying  
nonperturbative dynamics!**

# Nucleon- $\Delta$ - $\gamma$ transition



- **Magnetic dipole transition ( $G_M^*$ ) dominant:** quark spin flip (s wave). “Core + 25% pion cloud”
- **Electric & Coulomb quadrupole ratios** small & negative, encode deformation. Reproduced without pion cloud: **OAM from p waves!**

GE, Nicmorus, PRD 85 (2012)




# Compton scattering

## Nucleon polarizabilities:

### ChPT & dispersion relations

Hagelstein, Miskimen, Pascalutsa, PPNP 88 (2016)



In total: polarizabilities  $\approx$

Quark-level effects  $\leftrightarrow$  Baldin sum rule

+ nucleon resonances (mostly  $\Delta$ )

+ pion cloud (at low  $\eta_+$ )?

## First DSE results:

GE, FBS 57 (2016)

- Quark Compton vertex (Born + 1PI) calculated, added  $\Delta$  exchange

- compared to DRs


Pasquini et al., EPJ A11 (2001),

Downie & Fonvielle, EPJ ST 198 (2011)


- $\alpha_E$  dominated by handbag,  $\beta_M$  by  $\Delta$  contribution

$\Rightarrow$  large “QCD background”!

$\alpha_E + \beta_M$  [ $10^{-4}$  fm $^3$ ]




$\beta_M$  [ $10^{-4}$  fm $^3$ ]




# Tetraquarks in charm region?

- Can we **distinguish** different tetraquark configurations?




- Four quarks** dynamically rearrange themselves into  $dq\bar{d}\bar{q}$ , molecule, hadroquarkonium; strengths determined by four-body BSE:



# Muon g-2

- **Muon anomalous magnetic moment:**  
total SM prediction deviates from exp. by  $\sim 3\sigma$

Jegerlehner, Nyffeler,  
Phys. Rept. 477 (2009)



$$= ie \bar{u}(p') \left[ F_1(q^2) \gamma^\mu - F_2(q^2) \frac{\sigma^{\mu\nu} q_\nu}{2m} \right] u(p)$$

- Theory uncertainty dominated by **QCD**:  
Is QCD contribution under control?



Hadronic  
vacuum  
polarization




Hadronic  
light-by-light  
scattering

$a_\mu [10^{-10}]$

<b>Exp:</b>	11 659 208.9	(6.3)
<b>QED:</b>	11 658 471.9	(0.0)
<b>EW:</b>	15.3	(0.2)
<b>Hadronic:</b>		
• VP (LO+HO)	685.1	(4.3)
• <b>LBL</b>	<b>10.5</b>	<b>(2.6) ?</b>
<b>SM:</b>	11 659 182.8	(4.9)
<b>Diff:</b>	26.1	(8.0)

- **LbL amplitude:** ENJL & MD model results

Bijnens 1995, Hakayawa 1995, Knecht 2002, Melnikov 2004, Prades 2009, Jegerlehner 2009, Pauk 2014




$2$        $8 \dots 11$        $-1$        $2$        $-2$        $(\times 10^{-10})$

# Muon g-2

- **Muon anomalous magnetic moment:**  
total SM prediction deviates from exp. by  $\sim 3\sigma$

Jegerlehner, Nyffeler,  
Phys. Rept. 477 (2009)



$$= ie \bar{u}(p') \left[ F_1(q^2) \gamma^\mu - F_2(q^2) \frac{\sigma^{\mu\nu} q_\nu}{2m} \right] u(p)$$

- Theory uncertainty dominated by **QCD**:  
Is QCD contribution under control?




Hadronic  
vacuum  
polarization



Hadronic  
light-by-light  
scattering

- **LbL amplitude** at quark level, derived from **gauge invariance**:

GE, Fischer, PRD 85 (2012), Goecke, Fischer, Williams, PRD 87 (2013)



- **no double-counting, gauge invariant!**
- need to understand **structure of amplitude**

GE, Fischer, Heupel, PRD 92 (2015)

$a_\mu [10^{-10}]$

<b>Exp:</b>	11 659 208.9	(6.3)
<b>QED:</b>	11 658 471.9	(0.0)
<b>EW:</b>	15.3	(0.2)
<b>Hadronic:</b>		
• VP (LO+HO)	685.1	(4.3)
• <b>LBL</b>	<b>10.5</b>	<b>(2.6) ?</b>
<b>SM:</b>	11 659 182.8	(4.9)
<b>Diff:</b>	26.1	(8.0)