

Thermal ground state in deconfining Yang-Mills thermodynamics

Ralf Hofmann

ITP Universität Heidelberg

*2nd Winter Workshop on Non-Perturbative Quantum Field Theory
5-7 October 2011
Sophia-Antipolis (France)*

7 October 2011



Table of Contents I

Euclidean vs. real-time formulation of finite-temperature field theory

Periodicity, Matsubara sums, etc.

Results of perturbative approach to Yang-Mills in Euclidean formulation

Selfdual gauge fields in $SU(2)$ with $|k| = 1$

Trivial-holonomy calorons

Nontrivial holonomy: Static magnetic dipoles

Coarse graining, BPS saturation \Rightarrow inert field ϕ

Principles constraining the construction

Relevant limit in Diakonov's semiclassical approximation?

Kernel of a differential operator D

$D = \partial_\tau^2 + \left(\frac{2\pi}{\beta}\right)^2$ and potential for ϕ

Effective action

Coupling $k = 1$ to $k = 0$ effectively

Tree-level ground-state estimate

Winding to unitary gauge: \mathbf{Z}_2 degeneracy

Mass spectrum; outlook resummed radiative corrections

Table of Contents II

T dependence of e

Partition function, Legendre transformation, and evolution equation

Fixed points, attractor, and critical temperature

Prediction of P and ρ

Summary and outlook

Radiative corrections

Loop expansions

Stable but unresolved monopoles

The two other phases

Some remarks on physics implications

Euclidean finite-temperature field theory

- ▶ representation of partition function Z (real scalar ϕ) invented by Schwinger, Feynman 1950s, see e.g. [M. Le Bellac “Thermal Field Theory”]

$$\begin{aligned} Z &= \text{Tr} e^{-\beta H} = \mathcal{N} \int_{\phi(\mathbf{x},0)=\phi_\alpha(\mathbf{x})}^{\phi(\mathbf{x},\beta)=\phi_\alpha(\mathbf{x})} \prod_{\mathbf{x},\tau'} d\phi(\mathbf{x},\tau') \times \\ &\quad \exp \left[- \int_0^\beta d\tau'' \int d^3y \left(\frac{1}{2} \partial_{\tau''} \phi \partial_{\tau''} \phi + \frac{1}{2} \nabla \phi \cdot \nabla \phi + V(\phi) \right) \right] \\ &\equiv \mathcal{N} \int_{\phi(\mathbf{x},0)=\phi_\alpha(\mathbf{x})}^{\phi(\mathbf{x},\beta)=\phi_\alpha(\mathbf{x})} \prod_{\mathbf{x},\tau'} d\phi(\mathbf{x},\tau') \exp \left[- \int_0^\beta d\tau'' \int d^3y \mathcal{L}_E \right], \end{aligned}$$

where $\beta \equiv 1/T$.

- ▶ in gauge theory: admissible changes of gauge respect periodicity of A_μ
- ▶ in gauge-theory PT: additional gauge fixing required (Faddeev-Popov or better)

Euclidean finite-temperature field theory

- ▶ loop expansion of N -point functions in momentum space, propagator \bar{D}

$$\bar{D}(\mathbf{p}, \omega_n) = \frac{1}{\omega_n^2 + \mathbf{p}^2 + m^2},$$

where $\omega_n \equiv 2\pi nT$ ($n \in \mathbf{Z}$) n th Matsubara frequency.

- ▶ re-expressing (but not changing the contour for τ'' integration in Euclid. action) summation over n and integration over \mathbf{p} , $\sum_n \int d^3 p$, by Cauchy's integral theorem \Rightarrow

$$-\frac{1}{\omega_n^2 + \mathbf{p}^2 + m^2} \longrightarrow \frac{i}{p^2 - m^2} + \delta(p^2 - m^2) \frac{2\pi}{e^{\beta|p_0|} - 1},$$

where $\sum_n \int d^3 p \longrightarrow \int d^4 p$.

Real-time interpretation of loop integrals


Remarks:

- ▶ A more elaborate τ'' integration contour in the action was considered in [Umezawa, Matsumoto, and Tachiki (1982), Niemi and Semenoff (1984)]. This doubles real-time DOEs to avoid **pinch singularities** in PT.

- ▶ In Yang-Mills, where topological field configurations constructed for $0 \leq \tau'' \leq \beta$ (ground state!), such a change of contour for physics of propagating excitations is **inconsistent**.

Perturbative approach to pressure in Euclidean formulation

- ▶ in [Linde 1980] uselessness of PT after order g^6 pointed out (scale-separation argument for $g \ll 1$: momenta of order T (hard), gT (soft), and $g^2 T$ (ultrasoft); hard and soft OK; ultrasoft: weak screening of magnetic modes destroys perturbativity starting at g^6)
- ▶ SU(3) pressure in pure-YM PT



[Shuryak 1978, Kapusta 1979, Toimula 1983, Arnold and Zhai 1994, Zhai and Kastening 1994, Braaten and Nieto 1996, Kajantie 2003]

Trivial-holonomy calorons

- ▶ in singular gauge (winding number $|k| = 1$ is localized in a point) there is a **superposition principle** of instanton centers in **prepotential** Π [t Hooft (1976), Jackiw and Rebbi (1976)]:

$$\begin{aligned}\bar{A}_\mu^{+,a}(x) &= -\bar{\eta}_{\mu\nu}^a \partial_\nu \log \Pi, \\ \bar{A}_\mu^{-,a}(x) &= -\eta_{\mu\nu}^a \partial_\nu \log \Pi.\end{aligned}$$

- ▶ can be used to satisfy at $|k| = 1$ periodic b.c. in strip $(0 \leq \tau \leq \beta) \times \mathbf{R}^3$ [Harrington and Shepard (1978)]:

$$\begin{aligned}\Pi(\tau, \mathbf{x}; \rho, \beta, x_0) &= 1 + \sum_{l=-\infty}^{l=\infty} \frac{\rho^2}{(x - x_l)^2} \\ &= 1 + \frac{\pi \rho^2}{\beta r} \frac{\sinh\left(\frac{2\pi r}{\beta}\right)}{\cosh\left(\frac{2\pi r}{\beta}\right) - \cos\left(\frac{2\pi \tau}{\beta}\right)},\end{aligned}$$

where $r \equiv |\mathbf{x}|$.

Trivial-holonomy calorons, cntd.

- ▶ holonomy of $\bar{A}_\mu^{\pm,a}(x)$ at $r \rightarrow \infty$ trivial:

$$\prod_{r \rightarrow \infty} \left(1 + \frac{\pi \rho^2}{\beta r} \right) \Rightarrow \lim_{r \rightarrow \infty} \bar{A}_4^\pm \propto \lim_{r \rightarrow \infty} \frac{1}{r^2} = 0 \Rightarrow$$

$$\mathcal{P} \exp \left[i \int_0^\beta d\tau \bar{A}_4^\pm \right] = \mathbf{1}_2.$$

- ▶ Gaussian quantum weight [Gross, Pisarski, and Yaffe (1981)]:

$$S_{\text{eff}} = \frac{8\pi^2}{\bar{g}^2} + \frac{4}{3}\sigma^2 + 16 A(\sigma) \quad (\sigma \equiv \pi \frac{\rho}{\beta}),$$

$$A(\sigma) \rightarrow -\frac{1}{6} \log \sigma \quad (\sigma \rightarrow \infty) \quad A(\sigma) \rightarrow -\frac{\sigma^2}{36} \quad (\sigma \rightarrow 0).$$

Conclusion of **semiclassical approx.**:

Trivial-holonomy-caloron weight exponentially suppressed at high T .

Nontrivial holonomy: Static magnetic dipoles

- ▶ construction based on [Ward 1977, Atiyah and Ward 1977, ADHM 1978, Drinfeld and Manin 1978, Manton 1978, Adler 1978, Rossi 1979, Nahm 1980-1983]
- ▶ explicitly carried out in [Lee and Lu 1998, Kraan and Van Baal 1998]: $A_4(\tau, r \rightarrow \infty) = -iut^3 (0 \leq u \leq \frac{2\pi}{\beta})$.

exact cancellation
between A_4 -mediated
repulsion and
 A_i -mediated
attraction;

caloron radius ρ and
thus monopole-core
separation $D = \frac{\pi}{\beta} \rho^2$
increase from left to
right (T and
holonomy fixed)



action density of nontrivial-holonomy caloron with
 $k = 1$ plotted on 2D spatial slice

Nontrivial holonomy, cntd.

computation of functional determinant about nontrivial holonomy carried out in [Gross, Pisarski, and Yaffe (1981), Diakonov et al. 2004]

in (relevant) limit $\frac{D}{\beta} = \pi \left(\frac{\rho}{\beta} \right)^2 \gg 1$

conclusions:

- ▶ **total suppression** for nontrivial static holonomy in limit $V \rightarrow \infty$
- ▶ **attraction** of monop. and antimonop. for **small holonomy** ($0 \leq u \leq \frac{\pi}{\beta}(1 - \frac{1}{\sqrt{3}})$; $\frac{\pi}{\beta}(1 + \frac{1}{\sqrt{3}}) \leq u \leq 2 \frac{\pi}{\beta}$)
- ▶ **repulsion** of monop. and antimonop. for **large holonomy** ($\frac{\pi}{\beta}(1 - \frac{1}{\sqrt{3}}) \leq u \leq \frac{\pi}{\beta}(1 + \frac{1}{\sqrt{3}})$)
- ▶ **Instability** of classical configuration under quantum noise \Rightarrow **Nontrivial holonomy does not enter a priori estimate of thermal ground state!**

Inert field ϕ

Observations and principles constraining construction of ϕ :

- ▶ $F_{\mu\nu} = \pm \tilde{F}_{\mu\nu} \Rightarrow$ vanishing energy-momentum:

$$\Theta_{\mu\nu} = -2 \operatorname{tr} \left\{ \delta_{\mu\nu} \left(\mp \mathbf{E} \cdot \mathbf{B} \pm \frac{1}{4} (2\mathbf{E} \cdot \mathbf{B} + 2\mathbf{B} \cdot \mathbf{E}) \right) \right. \\ \left. \mp (\delta_{\mu 4} \delta_{\nu i} + \delta_{\mu i} \delta_{\nu 4}) (\mathbf{E} \times \mathbf{E})_i \right. \\ \left. \pm \delta_{\mu i} \delta_{\nu (j \neq i)} (E_i B_j - E_j B_i) \pm \delta_{\mu (j \neq i)} \delta_{\nu i} (E_j B_i - E_i B_j) \right\} \equiv 0.$$

- ▶ spatial isotropy and homogeneity of *effective* local field *not* associated with propagation of energy-momentum by *fundamental* gauge fields \Rightarrow **inert scalar** ϕ
- ▶ modulo admissible gauge transformations ϕ does not depend on time
- ▶ relevance of ϕ (BPS) by gauge-invariant coupling to coarse-grained $k = 0$ sector (perturbative renormalizability) \Rightarrow ϕ **adjoint** scalar

Inert field ϕ

Observations and principles constraining construction of ϕ , cntd:

- ▶ $F_{\mu\nu} \equiv \pm \tilde{F}_{\mu\nu} \Rightarrow$ any *local* “power” of $F_{\mu\nu}$ with an insertion of t^a **vanishes**
- ▶ **only trivial holonomy** in $F_{\mu\nu} \equiv \pm \tilde{F}_{\mu\nu}$ allowed
- ▶ $|\phi|$ is spacetime homogeneous \Rightarrow information on ϕ 's EOM is encoded in phase $\hat{\phi} \equiv \frac{\phi}{|\phi|}$
- ▶ definition of possible phases $\{\hat{\phi}\}$: due to BPS of A_μ^\pm **no explicit T dependence, flat measure** for admissible **integration over moduli** (excluding temporal shifts and global gauge rotations), Wilson lines between spatial points along **straight lines**

Inert field ϕ

Unique definition of $\{\hat{\phi}\}$ [Herbst and Hofmann 2004]:

$$\{\hat{\phi}^a\} \equiv \sum_{\pm} \text{tr} \int d^3x \int d\rho t^a F_{\mu\nu}(\tau, \mathbf{0}) \{(\tau, \mathbf{0}), (\tau, \mathbf{x})\} \\ \times F_{\mu\nu}(\tau, \mathbf{x}) \{(\tau, \mathbf{x}), (\tau, \mathbf{0})\} ,$$

where

$$\{(\tau, \mathbf{0}), (\tau, \mathbf{x})\} \equiv \mathcal{P} \exp \left[i \int_{(\tau, \mathbf{0})}^{(\tau, \mathbf{x})} dz_{\mu} A_{\mu}(z) \right] ,$$

$$\{(\tau, \mathbf{x}), (\tau, \mathbf{0})\} \equiv \{(\tau, \mathbf{0}), (\tau, \mathbf{x})\}^{\dagger} ,$$

and sum is over **Harrington-Shepard** (trivial-holonomy) caloron and anticaloron of scale ρ .

Higher n -point functions, higher topol. charge k ? **No.**

(Would introduce mass dimension $d = 3 - n - m$ of object, $m > 1$ number of dimension-length caloron moduli at $k > 1$, but d needs to vanish.)

Inert field ϕ

Some observations, conventions:


- ▶ $\hat{\phi}$ indeed transforms as an adjoint scalar:

$$\hat{\phi}^a(\tau) \rightarrow R^{ab}(\tau)\hat{\phi}^b(\tau),$$


where R^{ab} is τ dependent matrix of adjoint rep.

$$R^{ab}(\tau)t^b = \Omega^\dagger(\tau, \mathbf{0})t^a\Omega(\tau, \mathbf{0}).$$

- ▶ What about shift of spatial center $\mathbf{0} \rightarrow \mathbf{z}_\pm$?



(a)



(b)

Shift of center amounts to spatially *global* gauge rotation induced by the group element

$$\Omega_z^\pm = \{(\tau, \mathbf{0}), (\tau, \mathbf{z}_\pm)\}.$$

(a) graphical representation of **definition**

(b) only possible generalization to $\mathbf{z}_\pm \neq \mathbf{0}$

Inert field ϕ

Some observations, conventions, cntd:

- ▶ one has

$$\begin{aligned} \int_{(\tau, \mathbf{0})}^{(\tau, \mathbf{x})} dz_\mu A_\mu(z)|_\pm &= \pm \int_0^1 ds x_i A_i(\tau, s\mathbf{x}) \\ &= \pm t_b x_b \partial_\tau \int_0^1 ds \log \Pi(\tau, sr, \rho) \Rightarrow \end{aligned}$$

integrand in the exponent of $\{(\tau, \mathbf{0}), (\tau, \mathbf{x})\}_\pm$ varies along a fixed direction in $\mathfrak{su}(2)$ (a hedge hog); **Path-ordering can be ignored.**

- ▶ temporal shift freedom in A_μ^\pm : set $\tau_\pm = 0$ and re-instate later
- ▶ parity: $F_{\mu\nu}(\tau, \mathbf{x})_+ = F_{\mu\nu}(\tau, -\mathbf{x})_-$ and

$$\begin{aligned} \{(\tau, \mathbf{0}), (\tau, \mathbf{x})\}_+ &= (\{(\tau, \mathbf{x}), (\tau, \mathbf{0})\}_+)^{\dagger} = \{(\tau, \mathbf{0}), (\tau, -\mathbf{x})\}_- \\ &= (\{(\tau, -\mathbf{x}), (\tau, \mathbf{0})\}_-)^{\dagger} \Rightarrow \end{aligned}$$

– contribution to the integrand in **definition** obtained by $\mathbf{x} \rightarrow -\mathbf{x}$ in + contribution

Inert field ϕ

Some observations, conventions, cntd:

after tedious computation [Herbst and Hofmann 2004]

+ contribution to integrand in **definition** reads:

$$-i\beta^{-2} \frac{32\pi^4}{3} \frac{x^a}{r} \frac{\pi^2 \hat{\rho}^4 + \hat{\rho}^2(2 + \cos(2\pi\hat{\tau}))}{(2\pi^2 \hat{\rho}^2 + 1 - \cos(2\pi\hat{\tau}))^2} \times F[\hat{g}, \Pi],$$

where $\hat{\rho} \equiv \frac{\rho}{\beta}$, $\hat{r} \equiv \frac{r}{\beta}$, $\hat{\tau} \equiv \frac{\tau}{\beta}$, and functional F is

$$F[\hat{g}, \Pi] = 2 \cos(2\hat{g}) \left(2 \frac{[\partial_\tau \Pi][\partial_r \Pi]}{\Pi^2} - \frac{\partial_\tau \partial_r \Pi}{\Pi} \right) \\ + \sin(2\hat{g}) \left(2 \frac{[\partial_r \Pi]^2}{\Pi^2} - 2 \frac{[\partial_\tau \Pi]^2}{\Pi^2} + \frac{\partial_\tau^2 \Pi}{\Pi} - \frac{\partial_r^2 \Pi}{\Pi} \right),$$

and

$$\{(\tau, \mathbf{0}), (\tau, \mathbf{x})\}_\pm \equiv \cos \hat{g} \pm 2it_b \frac{x^b}{r} \sin \hat{g}.$$

One shows that \hat{g} saturates exponentially fast for $\hat{r} > 1$.

Inert field ϕ

discussion:


- ▶ angular integration would yield **zero** if radial integration was regular
- ▶ **but:** radial integration diverges logarithmically due to term $\frac{\partial_r^2 \Pi}{\Pi}$; this term arises from the **magnetic-magnetic** correlation (recall: no convergence in PT due to weakly screened magnetic sector!)
- ▶ zero \times infinity yields undetermined, multiplicative, and real constants Ξ_{\pm}
- ▶ without restriction of generality (global choice of gauge), angular integration regularized by defect azimuthal angle in 1-2 plane of $\mathfrak{su}(2)$ for both $+$ and $-$ contributions \Rightarrow **Members of $\{\hat{\phi}\}$ all move in hyperplane of $\mathfrak{su}(2)$!**
- ▶ re-instate $\tau \rightarrow \tau + \tau_{\pm} \Rightarrow$

Inert field ϕ

discussion, cntd:

result:

$$\{\hat{\phi}^a\} = \{\Xi_+(\delta^{a1} \cos \alpha_+ + \delta^{a2} \sin \alpha_+) \mathcal{A}(2\pi(\hat{\tau} + \hat{\tau}_+)) + \Xi_-(\delta^{a1} \cos \alpha_- + \delta^{a2} \sin \alpha_-) \mathcal{A}(2\pi(\hat{\tau} + \hat{\tau}_-))\}, \quad \text{where}$$



τ dependence of function $\mathcal{A}(\frac{2\pi\tau}{\beta})$;

saturation property (cutoff independence) for $\hat{\rho}$ integration.

ξ dependence of Ξ_{\pm}

$$\rho_{max} \equiv \xi\beta:$$

$$\int d\rho \rightarrow \int_0^{\zeta\beta} d\rho, \quad (\zeta > 0).$$

- ▶ $\Xi_{\pm} = 272 \zeta^3 \times \text{unknown, fixed real, } (\zeta > 10)$
- ▶ integral over ρ is strongly dominated by contributions just below upper limit
- ▶ semiclassical discussion of nontrivial-holonomy calorons in limit $\frac{D}{\beta} = \pi \left(\frac{\rho}{\beta}\right)^2 \gg 1$ [Diakonov et al. 2004] **is justified.**

Kernel of a differential operator D and potential for ϕ

- ▶ set $\{\hat{\phi}\}$ contains two real parameters for each “polarization”: Ξ_{\pm} and τ_{\pm} ; $\{\hat{\phi}\}$ is annihilated by **linear, second-order** differential operator $D = \partial_{\tau}^2 + \left(\frac{2\pi}{\beta}\right)^2 \Rightarrow$
 $\{\hat{\phi}\}$ coincides with **kernel** of D and determines D uniquely
- ▶ linearity \Rightarrow also $D\phi = 0$
- ▶ **but:** D depends on β explicitly, not allowed (BPS, caloron action given by topolog. charge)
- ▶ therefore seek potential $V(|\phi|^2)$ such that (Euclidean) action principle applied to

$$\mathcal{L}_{\phi} = \text{tr} \left((\partial_{\tau}\phi)^2 + V(\phi^2) \right) .$$

yields solutions annihilated by D , where \mathcal{L}_{ϕ} does not depend on β explicitly; demand that energy density $\Theta_{44} = 0$ on those solutions

Potential for and modulus of ϕ

- ▶ pick motion in 1-2 plane of $\mathfrak{su}(2)$ (gauge invariance $\Rightarrow V$ **central** potential \Rightarrow cons. angular momentum); ansatz:

$$\phi = 2 |\phi| t_1 \exp\left(\pm \frac{4\pi i}{\beta} t_3 \tau\right).$$

(circular motion in 1-2 plane, $|\phi|$ time independent!)

- ▶ apply E-L to $\mathcal{L}_\phi \Rightarrow$

$$\partial_\tau^2 \phi^a = \frac{\partial V(|\phi|^2)}{\partial |\phi|^2} \phi^a \quad (\text{in components}) \Leftrightarrow$$

$$\partial_\tau^2 \phi = \frac{\partial V(\phi^2)}{\partial \phi^2} \phi \quad (\text{in matrix form}).$$

- ▶ $\Theta_{44} = 0$ on ansatz $\phi \Rightarrow |\phi|^2 \left(\frac{2\pi}{\beta}\right)^2 - V(|\phi|^2) = 0$ but also:

$$\partial_\tau^2 \phi + \left(\frac{2\pi}{\beta}\right)^2 \phi = 0 \Rightarrow$$

$$\frac{\partial V(|\phi|^2)}{\partial |\phi|^2} = -\frac{V(|\phi|^2)}{|\phi|^2}.$$

Potential for and modulus of ϕ , cntd

$$\blacktriangleright \Rightarrow V(|\phi|^2) = \frac{\Lambda^6}{|\phi|^2}$$

where Λ integration constant of mass dim. unity.

$$\blacktriangleright \Rightarrow |\phi| = \sqrt{\frac{\Lambda^3 \beta}{2\pi}} \text{ (power-like decay of field } \phi \text{ with increasing } T)$$

*The field ϕ describes coarse-grained effect of **noninteracting** trivial-holonomy calorons and anticalorons. It does not propagate, and its modulus $|\phi|$ sets the scale of off-shellness down to which quantum fluctuations, arising from the sector $k = 0$, must be considered “integrated out” in full effective theory.*

\blacktriangleright Indeed: cutting off ρ and r integrations at $|\phi|^{-1}$, τ dependence of $\mathcal{A}(\frac{2\pi\tau}{\beta})$ is perfect sine

(Error at level smaller than 10^{-22} if knowledge about

$T_c = \frac{\lambda_c \Lambda}{2\pi}$ with $\lambda_c = 13.87$ is used, later.)

BPS equation for ϕ

In addition to E-L equation ϕ satisfies **first-order**, BPS equation:


$$\partial_\tau \phi = \pm 2i \Lambda^3 t_3 \phi^{-1} = \pm i V^{1/2}(\phi).$$

*Because ϕ satisfies both, second-order E-L and first-order BPS equation, usual shift ambiguity in ground-state energy density, as allowed by E-L equation, **absent** in $SU(2)$ Yang-Mills thermodynamics.*

Effective action for deconfining phase

Coupling the coarse-grained $k = 0$ sector to ϕ , following constraints:

- ▶ perturbative renormalizability
 ['t Hooft, Veltman, Lee, and Zinn-Justin 1971-1973]
 \Rightarrow form invariance of action for effective $k = 0$ gauge field a_μ from integrating fundamental $k = 0$ fluctuations only, no higher dim. ops. for a_μ only
- ▶ no energy-momentum transfer to $\phi \Rightarrow$ absence of higher dim. ops. involving a_μ **and** ϕ
- ▶ gauge invariance $\Rightarrow \partial_\mu \phi \rightarrow D_\mu \phi \equiv \partial_\mu \phi - ie[a_\mu, \phi]$ (**effective** coupling); no momentum transfer to ϕ if (unitary gauge $\phi = 2|\phi| t_3$) massive 1,2 modes propagate on-shell only



Effective action and ground-state estimate

unique effective action density:

$$\mathcal{L}_{\text{eff}}[a_\mu] = \text{tr} \left(\frac{1}{2} G_{\mu\nu} G_{\mu\nu} + (D_\mu \phi)^2 + \frac{\Lambda^6}{\phi^2} \right),$$

$$\text{where } G_{\mu\nu} = \partial_\mu a_\nu - \partial_\nu a_\mu - ie[a_\mu, a_\nu] \equiv G_{\mu\nu}^a t_a$$

ground-state estimate:

- ▶ E-L EOM from $\mathcal{L}_{\text{eff}}[a_\mu]$

$$D_\mu G_{\mu\nu} = ie[\phi, D_\nu \phi].$$

- ▶ solved by zero-curvature (pure-gauge) config. a_μ^{gs} :

$$a_\mu^{\text{gs}} = \mp \delta_{\mu 4} \frac{2\pi}{e\beta} t_3 \quad (D_\nu \phi \equiv G_{\mu\nu} \equiv 0) \Rightarrow$$

$$\rho^{\text{gs}} = -P^{\text{gs}} = 4\pi\Lambda^3 T.$$

*Unresolvable interactions between $k = 0$ and $|k| = 1$ lifted ρ^{gs} from zero (BPS). EOS of a cosmological constant; pressure **negative**. (Short-lived, attracting magnetic (anti)monopoles by temporary shifts of (anti)caloron holonomies from trivial to small through absorption of hard plane-wave fluctuations.)*

Winding to unitary gauge: \mathbf{Z}_2 degeneracy


- ▶ consider gauge rotation $\tilde{\Omega}(\tau) = \Omega_{\text{gl}} Z(\tau) \Omega(\tau)$ where $\Omega(\tau) \equiv \exp[\pm 2\pi i \frac{\tau}{\beta} t_3]$, $Z(\tau) = \left(2\Theta(\tau - \frac{\beta}{2}) - 1\right) \mathbf{1}_2$, and $\Omega_{\text{gl}} = \exp[i\frac{\pi}{2} t_2]$
- ▶ $\tilde{\Omega}(\tau)$ transforms a_μ^{gs} to $a_\mu^{\text{gs}} \equiv 0$ and ϕ to $\phi = 2t^3|\phi|$
- ▶ $\tilde{\Omega}(\tau)$ is **admissible** because respects periodicity of δa_μ :

$$\begin{aligned} a_\mu &\rightarrow \tilde{\Omega}(a_\mu^{\text{gs}} + \delta a_\mu)\tilde{\Omega}^\dagger + \frac{i}{e}\tilde{\Omega}\partial_\mu\tilde{\Omega}^\dagger \\ &= \Omega_{\text{gl}} \left(\Omega(a_\mu^{\text{gs}} + \delta a_\mu)\Omega^\dagger + \frac{i}{e} \left(\Omega\partial_\mu\Omega^\dagger + Z\partial_\mu Z \right) \right) \Omega_{\text{gl}}^\dagger \\ &= \Omega_{\text{gl}} \left(\Omega\delta a_\mu\Omega^\dagger + \frac{2i}{e}\delta(\tau - \frac{\beta}{2})Z \right) \Omega_{\text{gl}}^\dagger = \Omega_{\text{gl}}\Omega \delta a_\mu (\Omega_{\text{gl}}\Omega)^\dagger. \end{aligned}$$

- ▶ $\tilde{\Omega}(\tau)$ transforms Polyakov loop from $-\mathbf{1}_2$ to $\mathbf{1}_2 \Rightarrow$
ground-state estimate is (electric) \mathbf{Z}_2 degenerate \Rightarrow
deconfining phase

Mass spectrum; outlook resummed radiative corrections

- ▶ computation in physical and completely fixed **unitary, Coulomb gauge** ($\phi = 2t^3|\phi|$, $\partial_i a_i^3 = 0$)
- ▶ mass spectrum: $m^2 \equiv m_1^2 = m_2^2 = 4e^2 \frac{\Lambda^3}{2\pi T}$, $m_3 = 0$.
- ▶ resummation of **polarization tensor of massless mode** as



⇒ small linear-in- T correction to tree-level ground-state estimate [Falquez, Hofmann, Baumbach 2010]

$$\begin{aligned} \text{tree-level:} & \quad \frac{\rho^{\text{gs}}}{T^4} = 3117.09 \lambda^{-3}, \\ \text{one-loop resummed:} & \quad \frac{\Delta\rho^{\text{gs}}}{T^4} = 3.95 \lambda^{-3}. \end{aligned}$$

- ▶ large hierarchy between loop orders (conjecture about **termination at finite irreducible order**, see second talk), so one-loop correction **practically exact**

T dependence of ϵ : selfconsistent thermal quasiparticles

P and ρ at one loop:

$$P(\lambda) = -\Lambda^4 \left\{ \frac{2\lambda^4}{(2\pi)^6} [2\bar{P}(0) + 6\bar{P}(2a)] + 2\lambda \right\},$$

$$\rho(\lambda) = \Lambda^4 \left\{ \frac{2\lambda^4}{(2\pi)^6} [2\bar{\rho}(0) + 6\bar{\rho}(2a)] + 2\lambda \right\},$$

where

$$\bar{P}(y) \equiv \int_0^\infty dx x^2 \log \left[1 - \exp(-\sqrt{x^2 + y^2}) \right],$$

$$\bar{\rho}(y) \equiv \int_0^\infty dx x^2 \frac{\sqrt{x^2 + y^2}}{\exp(\sqrt{x^2 + y^2}) - 1},$$

and $a \equiv \frac{m}{2T} = 2\pi e\lambda^{-3/2}$. For later use introduce function $D(2a)$ as

$$\partial_{y^2} \bar{P} \Big|_{y=2a} = -\frac{1}{4\pi^2} \int_0^\infty dx \frac{x^2}{\sqrt{x^2 + (2a)^2}} \frac{1}{e^{\sqrt{x^2 + (2a)^2}} - 1} \equiv -\frac{1}{4\pi^2} D(2a).$$

Legendre transformation and evolution equation

- ▶ for $m(T)$ to respect Legendre trafo (fundamental partition function) between P and $\rho \Leftrightarrow \partial_m P = 0$
- ▶ \Rightarrow first-order **evolution equation**

$$\partial_a \lambda = -\frac{24\lambda^4 a}{(2\pi)^6} \frac{D(2a)}{1 + \frac{24\lambda^3 a^2}{(2\pi)^6} D(2a)}.$$

or

$$1 = -\frac{24\lambda^3}{(2\pi)^6} \left(\lambda \frac{da}{d\lambda} + a \right) a D(2a).$$

- ▶ \Rightarrow dependence $a(\lambda)$ monotonic decreasing
 \Rightarrow for $\lambda \gg 1$ a must fall below unity
- ▶ **fixed points of evolution equation:**

repulsive at $a = 0$ ($\lambda \rightarrow \infty$)

attractive at $a = \infty$ ($\lambda = \lambda_c$)

Solution to evolution equation

- ▶ $a \ll 1$ [Dolan, Jackiw 1974] $\Rightarrow 1 = -\frac{\lambda^3}{(2\pi)^4} \left(\lambda \frac{da}{d\lambda} + a \right) a$;
solution ($a(\lambda_i) = a_i \ll 1$):


$$a(\lambda) = 4\sqrt{2}\pi^2 \lambda^{-3/2} \left(1 - \frac{\lambda}{\lambda_i} \left[1 - \frac{a_i^2 \lambda_i^3}{32\pi^4} \right] \right)^{1/2}.$$

\Rightarrow attractor $a(\lambda) = 4\sqrt{2}\pi^2 \lambda^{-3/2}$ as long as $a \ll 1$



$\Rightarrow e = \sqrt{8\pi}$ as long as $a \ll 1$ (**amusingly**: $S = \frac{8\pi^2}{e^2} = 1$)

(geometric interpretation of \hbar in terms of caloron winding number)

- ▶ full solution for $e(\lambda) \Rightarrow \lambda_c = 13.87$:



T dependence of P and ρ



- ▶ notice **negativity** of P shortly above λ_c
- ▶ relative correction to one-loop quasiparticle P and ρ by radiative effects: $< 1\%$, see second talk

Summary

Summary:

- ▶ brief motivation why nonperturbative approach to YMTD necessary: mass generation, poor convergence of pert. orders
- ▶ mini review on calorons: trivial vs. nontrivial holonomy for $|k| = 1$ plus semiclassical approx.
- ▶ construction of thermal ground-state estimate: inert field ϕ ; BPS and E-L; potential
- ▶ discussion of constraints on effective action: pert. renormalizability plus inertness of $\phi \Rightarrow$ unique answer
- ▶ full ground-state estimate, deconfining nature, tree-level quasiparticles
- ▶ evolution of effective coupling
- ▶ T dependence pressure and energy density

Outlook

- ▶ running of fundamental coupling: trace anomaly
- ▶ radiative corrections: polarization tensor of massless mode
- ▶ radiative corrections: stable but unresolvable monopoles
- ▶ radiative corrections: two-loop and three-loop cases
- ▶ radiative corrections: loop expansion of pressure, conjecture on termination at finite irreducible order
- ▶ two other phases:
 - ▶ **preconfining** (thermal ground state: condensate of massless monopoles and antimonopoles)
 - ▶ **confining** (ground state of zero energy density: condensate of single, round-point like center-vortex loops)

Physics

Some physics implications:

(i) mechanism for ew SB (LHC: not much of a Higgs signal so far)

(ii) postulate: $SU(2)$ (10^{-4} eV) describes photon **propagation**

⇒ black-body spectral anomaly at $T \sim 5 - 20$ K and low frequencies

(cold H1 clouds, large-angle anomalies in TT of CMB, UEGE)

⇒ Planck-scale axion plus such an $SU(2)$ yield **Dark Energy**

Thank you.