## PROBLEMS FOR QUANTUM FIELD THEORY 2 12. Sheet

Suggested reading before solving these problems: Chapter 6.2-6.4 in the script and/or chapter 6, 7, 11.1 in *Rothe: Lattice Gauge Theories*.

Problem 1: Strong coupling expansion in Lattice QCD

Consider SU(3) Yang-Mills theory on the lattice. The action is given by

$$S = -\beta \sum_{P} S_{P}$$

where  $\beta = \frac{6}{g_0^2}$  and

$$S_P = \frac{1}{6} \text{Tr}(U_P + U_P^{\dagger})$$

is the contribution associated with a plaquette P. The sum goes over all distinct plaquettes.  $U_P$  is the plaquette variable and can be expressed in terms of the link variables around the plaquette P as

$$U_P = U_{\mu\nu}(n) = U_{\mu}(n)U_{\nu}(n+\hat{\mu})U_{\mu}^{\dagger}(n+\hat{\nu})U_{\nu}^{\dagger}(n).$$

The Wlison loop with spatial extend  $\hat{R}$  and temporal extend  $\hat{T}$  (in units of the lattice spacing a) can be expressed in terms of the link variables by the path ordered product

$$W_C[U] = \operatorname{Tr} \prod U_l$$

of link variables around the loop.

Consider now the expectation value

$$\langle W_C[U] \rangle = \frac{\int DU W_C[U] e^{-S[U]}}{\int DU e^{-S[U]}}.$$

Expand the exponential in powers of the coupling  $\beta$ 

$$e^{-S[U]} = e^{\beta \sum_P S_P} = \prod_P \left[ \sum_n \frac{\beta^n}{n!} (S_P)^n \right]$$

and use the following integration rules for the integrals over the link variables  $U \in SU(3)$ 

$$\int dU \ U^{ab} = 0,$$

$$\int dU \ U^{ab} U^{cd} = 0,$$

$$\int dU \ U^{ab} (U^{\dagger})^{cd} = \frac{1}{3} \delta_{ad} \delta_{bc},$$

$$\int dU \ 1 = 1,$$

to show

$$\langle W_C[U] \rangle \sim \left(\frac{\beta}{c}\right)^{\hat{R}\hat{T}}$$

with some coefficient c.

The Wilson loop behaves for large  $\hat{T}$  as

$$\langle W_C[U] \rangle \stackrel{\hat{T} \to \infty}{\to} F(\hat{R}) e^{-\hat{V}(\hat{R})\hat{T}}$$

so that you can deduce that the potential is linear in  $\hat{R}$ :

$$\hat{V}(\hat{R}) = \hat{\sigma}\hat{R},$$

with  $\hat{\sigma} = -\ln(\beta/c)$  the string tension.