

---

# Quantum Field Theory 2 – Problem set 2

Lectures: Jan Pawłowski

j.pawlowski@thphys.uni-heidelberg.de

Tutorials: Eduardo Grossi

e.grossi@thphys.uni-heidelberg.de

Institut für Theoretische Physik, Uni Heidelberg

due date: 25 April 2017

---

## Problem 1: Partition function for imaginary temperature

Consider the canonical partition function

$$Z = \text{Tr} e^{-\beta H} = \sum_n \langle n | e^{-\beta H} | n \rangle$$

where  $|n\rangle$  is an arbitrary basis and  $\beta = 1/T$ . Show that this can be written as

$$Z = \int dq_0 \langle q_0 | e^{-\beta H} | q_0 \rangle$$

where  $|q_0\rangle$  labels eigenstates of the position operator. Let us assume for the moment that the temperature is imaginary so that we can write

$$\beta = it$$

with  $t > 0$ . Use the path integral formula for the transition amplitude

$$\langle q_f | e^{-iHt} | q_i \rangle = \int_{q(0)=q_i, q(t)=q_f} Dq e^{iS[q]},$$

$$S[q] = \int_0^t dt' L(q, \dot{q}, t')$$

to show that  $Z$  can be written as a functional integral over periodic functions  $q(t')$  with  $q(t) = q(0)$ .