
Quantum Field Theory 2 – Tutorial 4

Lectures: Jan Pawłowski

J.Pawlowski@thphys.uni-heidelberg.de

Tutorials: Aleksandr Mikheev

A.Mikheev@thphys.uni-heidelberg.de

Institut für Theoretische Physik, Uni Heidelberg

tutorial date: week of 08.05.2023

Problem 1: Legendre transform

Consider a real function $f(x)$ of a real variable x . The Legendre transform of $f(x)$ is defined by

$$g(y) = \sup_x (xy - f(x)). \quad (1)$$

How can one construct the function $g(y)$ graphically? Do this for $f(x) = \lambda x^4 - \mu x^2$. What is the Legendre transform of $g(y)$?

Problem 2: Effective action for Gaussian theory

Consider a probabilistic theory for a N -dimensional vector \mathbf{x} . The partition function is given by

$$Z(\mathbf{J}) = e^{W(\mathbf{j})} = \int d^N \tilde{x} e^{-S(\tilde{\mathbf{x}}) + J_n \tilde{x}_n}. \quad (2)$$

The effective action is defined as the Legendre transform of the Schwinger function $W(\mathbf{J})$,

$$\Gamma(\mathbf{x}) = \sup_{\mathbf{J}} (J_n x_n - W(\mathbf{J})). \quad (3)$$

Calculate $\Gamma(\mathbf{x})$ for the Gaussian theory $S(\mathbf{x}) = \frac{1}{2} P_{mn} x_m x_n$.