

---

# Quantum Field Theory 1 – Problem set 11

Lectures: Jan Pawłowski  
Tutorials: Michael Scherer

pawłowski@thphys.uni-heidelberg.de  
scherer@thphys.uni-heidelberg.de

Institut für Theoretische Physik, Uni Heidelberg

due date: 18 January 2017

---

## Problem 1: Dimensional regularization

Dimensional regularization is a method to regularize expressions that formally diverge in  $d = 3 + 1$  dimensions. One takes  $d$  to be a real number, for example  $d = 4 - 2\epsilon$ . A typical example is a (Euclidean or Wick-rotated) integral of the form

$$\int \frac{d^d q}{(2\pi)^d} \frac{1}{(q^2 + m^2)^n} = \frac{1}{2} \int \frac{d\Omega_d}{(2\pi)^d} 2 \int dq \frac{q^{d-1}}{(q^2 + m^2)^n}. \quad (1)$$

- a) The first integral in Eq. (1) contains the area of a unit sphere in  $d$  dimensions. By using  $(\sqrt{\pi})^d = \left( \int dx e^{-x^2} \right)^d = \int d^d x e^{-\sum_{i=1}^d x_i^2}$ , show that

$$\frac{1}{2} \int \frac{d\Omega_d}{(2\pi)^d} = \frac{1}{(4\pi)^{d/2} \Gamma(d/2)},$$

with the Gamma function  $\Gamma(x) = \int_0^\infty dt t^{x-1} e^{-t}$ .

- b) By substituting  $x = m^2/(q^2 + m^2)$  show that the remaining integral in Eq. (1) can be written as

$$2 \int dq \frac{q^{d-1}}{(q^2 + m^2)^n} = \left( \frac{1}{m^2} \right)^{n-\frac{d}{2}} \int_0^1 dx x^{n-\frac{d}{2}-1} (1-x)^{\frac{d}{2}-1}.$$

Use now the formula  $\int_0^1 dx x^{\alpha-1} (1-x)^{\beta-1} = \Gamma(\alpha)\Gamma(\beta)/\Gamma(\alpha+\beta)$  to show

$$\int \frac{d^d q}{(2\pi)^d} \frac{1}{(q^2 + m^2)^n} = \frac{1}{(4\pi)^{d/2} \Gamma(\frac{d}{2})} \left( \frac{1}{m^2} \right)^{n-\frac{d}{2}} \frac{\Gamma(n-\frac{d}{2})\Gamma(\frac{d}{2})}{\Gamma(n)}. \quad (2)$$

- c) Derive from Eq. (2)

$$\int \frac{d^d q}{(2\pi)^d} \frac{1}{(q^2 + m^2 + 2q \cdot p)^n} = \frac{\Gamma(n-\frac{d}{2})}{(4\pi)^{d/2} \Gamma(n)} \frac{1}{(m^2 - p^2)^{n-\frac{d}{2}}},$$

and use this together with properties of the Gamma function to show


$$\int \frac{d^d q}{(2\pi)^d} \frac{q^\mu}{(q^2 + m^2 + 2q \cdot p)^n} = - \frac{\Gamma(n-\frac{d}{2})}{(4\pi)^{d/2} \Gamma(n)} \frac{p^\mu}{(m^2 - p^2)^{n-\frac{d}{2}}}$$

and

$$\int \frac{d^d q}{(2\pi)^d} \frac{q^\mu q^\nu}{(q^2 + m^2 + 2q \cdot p)^n} = \frac{1}{(4\pi)^{d/2} \Gamma(n)} \left[ p^\mu p^\nu \frac{\Gamma(n-\frac{d}{2})}{(m^2 - p^2)^{n-\frac{d}{2}}} + \frac{1}{2} \delta^{\mu\nu} \frac{\Gamma(n-\frac{d}{2}-1)}{(m^2 - p^2)^{n-\frac{d}{2}-1}} \right].$$

## Problem 2: Vacuum polarization of QED

In this problem we calculate a one-loop contribution to the photon propagator



- a) Use the Feynman rules of QED to show that this amplitude reads

$$i\mathcal{M} = \epsilon_\mu^*(p) (i\Pi^{\mu\nu}(p)) \epsilon_\nu(p)$$

with

$$i\Pi^{\mu\nu}(p) = -(-ie)^2 \int \frac{d^4q}{(2\pi)^4} \text{tr} \left[ \gamma^\mu \frac{i(\not{q} + m)}{q^2 - m^2} \gamma^\nu \frac{i(\not{q} + \not{p} + m)}{(q+p)^2 - m^2} \right].$$

- b) Use now the trace identity

$$\text{tr} \gamma^\mu \gamma^\nu \gamma^\sigma \gamma^\rho = 4(\eta^{\mu\nu} \eta^{\sigma\rho} + \eta^{\mu\rho} \eta^{\nu\sigma} - \eta^{\mu\sigma} \eta^{\nu\rho}),$$

the Feynman trick

$$\frac{1}{AB} = \int d\alpha \frac{1}{(\alpha A + (1-\alpha)B)^2},$$

an appropriate shift in the integration variable and Wick rotation to show

$$i\Pi^{\mu\nu}(p) = -4ie^2 \int_0^1 d\alpha \int \frac{d^d q}{(2\pi)^d} \frac{-\frac{2}{d}\eta^{\mu\nu} q^2 + \eta^{\mu\nu} q^2 - 2\alpha(1-\alpha)p^\mu p^\nu + \eta^{\mu\nu}(m^2 + \alpha(1-\alpha)p^2)}{(q^2 + \Delta)^2}$$

with  $\Delta = m^2 - \alpha(1-\alpha)p^2$ .

- c) You can now use the results of problem 1 as well as the expansion ( $d = 4 - 2\epsilon$ )

$$\Gamma(2 - \frac{d}{2}) = \Gamma(\epsilon) = \frac{1}{\epsilon} - \gamma + \mathcal{O}(\epsilon),$$

with  $\gamma \approx 0.5772$  the Euler-Mascheroni constant to show that  $\Pi^{\mu\nu}(p)$  is of the form

$$i\Pi^{\mu\nu}(p) = (p^2 \eta^{\mu\nu} - p^\mu p^\nu) i\Pi(p^2)$$

with

$$\Pi(p^2) = -\frac{8e^2}{(4\pi)^{d/2}} \int_0^1 d\alpha \alpha(1-\alpha) \frac{\Gamma(2 - \frac{d}{2})}{\Delta^{2 - \frac{d}{2}}} \xrightarrow{d \rightarrow 4} -\frac{e^2}{2\pi^2} \int_0^1 d\alpha \alpha(1-\alpha) \left( \frac{1}{\epsilon} - \ln \Delta - \gamma \right).$$