

---

# Quantum Field Theory 1 – Tutorial 1

Lectures: Jan Pawłowski

pawłowski@thphys.uni-heidelberg.de

Tutorials: Michael Scherer

scherer@thphys.uni-heidelberg.de

Institut für Theoretische Physik, Uni Heidelberg

tutorial date: 19 October 2016

---

## Problem 1: Natural units

In particle physics one often chooses units such that  $\hbar = c = 1$  and measures energies in MeV. Complete the following translation table (The charge of a proton is  $e = 1.6 \cdot 10^{-19} C$ ):

|   | SI units   | Natural units |
|---|--|---------------|
| $c$   | $3 \cdot 10^8 \text{ m/s}$                       | 1             |
| $\hbar$                                       | $1.05 \cdot 10^{-34} \text{ Js}$                 | 1             |
| $m_e$   | $9.1 \cdot 10^{-31} \text{ kg}$                  |               |
| $m_p$   |  | 938.3 MeV     |
| $\alpha = \frac{e^2}{4\pi\epsilon_0\hbar c}$  |  |               |
| $a_0 = \frac{4\pi\epsilon_0\hbar^2}{m_e e^2}$ | $0.53 \cdot 10^{-10} \text{ m}$                  |               |
| $G$   | $6.67 \cdot 10^{-11} \text{ Nm}^2/(\text{kg}^2)$ |               |

## Problem 2: Upper and lower indices

In special relativity one distinguishes between upper or contra-variant indices (as e.g.  $x^\mu$ ) and lower or co-variant indices (e.g.  $x_\mu$ ). The metric tensor

$$(\eta_{\mu\nu}) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

is used to raise or lower indices:

$$x_\mu = \eta_{\mu\nu} x^\nu, \quad x^\mu = \eta^{\mu\nu} x_\nu.$$

For  $x^\mu = (x^0, \mathbf{x})$  and  $p^\mu = (p^0, \mathbf{p})$  calculate

$$x_\mu, \quad p \cdot x = p_\mu x^\mu$$

and show that

$$\eta^\mu{}_\nu = \delta^\mu{}_\nu, \quad (\eta^{\mu\nu}) = (\eta_{\mu\nu}).$$

### Problem 3: Poincaré group

Under a Poincaré transformation  $(\Lambda, a)$  a coordinate vector  $x^\mu$  transforms as

$$x^\mu \rightarrow x'^\mu = \Lambda^\mu{}_\nu x^\nu + a^\mu.$$

Poincaré transformations leave the scalar product of differences of coordinate vectors,  $(x - y) \cdot (x - y)$ , invariant. Hence the matrix  $\Lambda$  satisfies

$$\Lambda^\rho{}_\mu \Lambda^\sigma{}_\nu \eta^{\mu\nu} = \eta^{\rho\sigma}. \quad (1)$$

a) Show that

$$(\Lambda^{-1})^\mu{}_\nu = \Lambda_\nu{}^\mu.$$

b) Consider the product of two Poincaré transformations

$$(\Lambda, a) = (\Lambda_1, a_1)(\Lambda_2, a_2).$$

Determine  $(\Lambda, a)$  and show that  $\Lambda$  satisfies Eq. (1).

c) Determine the inverse transformation

$$(\Lambda, a)^{-1}.$$

*Remark:* The properties shown here together with associativity

$$\left[ (\Lambda_1, a_1)(\Lambda_2, a_2) \right] (\Lambda_3, a_3) = (\Lambda_1, a_1) \left[ (\Lambda_2, a_2)(\Lambda_3, a_3) \right]$$

and the existence of a unit element

$$(\mathbb{1}, 0)$$

imply that the set of Poincaré transformations constitutes a *group*.