

---

# Quantum Field Theory 1 – Tutorial 11

Lectures: Jan Pawłowski

pawłowski@thphys.uni-heidelberg.de

Tutorials: Michael Scherer

scherer@thphys.uni-heidelberg.de

Institut für Theoretische Physik, Uni Heidelberg

tutorial date: 11 January 2017

---

## Problem 1: QED Lagrangian and gauge fixing

The pure gauge field Lagrangian  $\mathcal{L}_A = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$  gives an equation of motion of the form  $Q^{\nu\sigma}A_\sigma = 0$  with operator  $Q^{\nu\sigma} = \partial_\mu\partial^\mu\eta^{\nu\sigma} - \partial^\nu\partial^\sigma$ .

- a) Show that  $Q^{\nu\sigma}$  is not invertible as it annihilates any function of the form  $\partial_\sigma\alpha$ .

This prevents us from obtaining a sensible photon propagator directly from  $\mathcal{L}_A$  and is related to the redundancy of the gauge field,  $A_\mu \rightarrow A_\mu + e\partial_\mu\alpha$ . To remove the redundancy, we introduce a *gauge fixing* (cf. script Eq. (5.21), p. 138)

$$\partial_\mu A^\mu = 0. \tag{1}$$

- b) Show that for any given field  $A'_\mu$  we can always pick a representative configuration, which satisfies Eq. (1). Specify the required gauge transformation.
- c) Show that Lorenz gauge leaves a subspace of gauge transformations which satisfy Eq. (1). Which equation has to be satisfied by further gauge transformations?

A complete the gauge fixing can be achieved by adding appropriate boundary conditions. With the gauge fixing constraint, we can then write the gauge field Lagrangian as

$$\mathcal{L}_A = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{2\xi}(\partial_\mu A^\mu)^2, \tag{2}$$

and obtain the invertible operator  $\partial_\mu\partial^\mu\eta^{\nu\sigma} - \left(1 - \frac{1}{\xi}\right)\partial^\nu\partial^\sigma$ , cf. script Eq. (5.22), p. 139.