Quantum Field Theory 1 – Problem set 5

Lectures: Jörg Jäckel Jan Pawlowski Tutorials: Malo Tarpin Institut für Theoretische Physik, Uni Heidelberg

J.Jaeckel@thphys.uni-heidelberg.de J.Pawlowski@thphys.uni-heidelberg.de M.Tarpin@thphys.uni-heidelberg.de tutorial date: 11 November 2016

Suggested reading before solving these problems: Chapters 3.1 to 3.2 in the script and/or Chapters 4.2 to 4.3 of *Peskin & Schroeder*.

Problem 1: Time and normal ordering 1

Explain in a few words and formula

- a) What is a time ordered product and why do we need it?
- b) What is a normal ordered product and why is it useful?
- c) How can we transform a time ordered expression to a normal ordered one?

Problem 2: Time and normal ordering 2

- a) Show that the time ordered product $T(\phi(x_1)\phi(x_2))$ and the normal ordered product $: \phi(x_1)\phi(x_2) :$ are both symmetric under the interchange of x_1 and x_2 . Deduce that the Feynman propagator $D_F(x_1 x_2)$ has the same property.
- b) Check Wick's theorem for the case of three real scalar fields

$$T(\phi(x_1)\phi(x_2)\phi(x_3)) = :\phi(x_1)\phi(x_2)\phi(x_3):+\phi(x_1)D_F(x_2-x_3) +\phi(x_2)D_F(x_1-x_3)+\phi(x_3)D_F(x_1-x_2).$$

Problem 3: Three-point correlation function

Consider the following Lagrangian, involving two real scalar fields Φ and ϕ :

$$\mathcal{L} = \frac{1}{2}\partial_{\mu}\Phi\partial^{\mu}\Phi - \frac{1}{2}M^{2}\Phi^{2} + \frac{1}{2}\partial_{\mu}\phi\partial^{\mu}\phi - \frac{1}{2}m^{2}\phi^{2} - h\Phi\phi\phi.$$

The last term is an interaction term that allows a Φ particle to decay into two ϕ 's, provided that M > 2m. Assume that this is the case and calculate the $\mathcal{O}(h)$ contribution to the correlation function

 $\langle 0| T \Phi(x) \phi(y) \phi(z) e^{-i \int dt H_{\text{int}}(t)} |0\rangle.$