
Quantum Field Theory 1 – Tutorial 5

Lectures: Jan Pawłowski

J.Pawłowski@thphys.uni-heidelberg.de

Tutorials: Felipe Attanasio

F.Attanasio@thphys.uni-heidelberg.de

Institut für Theoretische Physik, Uni Heidelberg

tutorial date: week of 14.11.2022

Problem 1: Feynman propagator

(a) Calculate the integral

$$\int_{-\infty}^{\infty} \frac{e^{itx}}{x^2 + 1} dx$$

for $t > 0$ using the residue theorem.

(b) Consider the Feynman propagator for a scalar field

$$D_F(x - y) = \int \frac{d^4p}{(2\pi)^4} \frac{i}{p^2 - m^2 + i\epsilon} e^{-ip(x-y)}$$

Assuming $x_0 - y_0 > 0$ and using the residue theorem, show that this can be written as

$$D_F(x - y) = \int \frac{d^3p}{(2\pi)^3} \frac{1}{2E_p} e^{-iE_p(x_0 - y_0) + i\mathbf{p}(\mathbf{x} - \mathbf{y})}.$$

How does this change for $x_0 - y_0 < 0$?

Residue theorem: The contour integral of a function $f(z)$ around a closed, counter-clockwise path encircling a domain where $f(z)$ has a finite number of isolated singularities (poles at $z = z_i, i = 1, 2, \dots, n$) is

$$\oint dz f(z) = 2\pi i \sum_{i=1}^n \text{Res}(f, z_i)$$

where the residue of $f(z)$ at a simple pole z_i is $\text{Res}(f, z_i) = \lim_{z \rightarrow z_i} (z - z_i) f(z)$.