


Quantum fermions from classical bits



Fermions are Ising spins

- Fermionic occupation numbers $n = 0, 1$
- Classical bits
- Ising spins $s = \pm 1$
- Bit configurations = many body states of fermions


Cellular automaton

At each step :

- each bit configuration changes to a unique new bit configuration according to an updating rule
- for a fixed initial configuration : classical deterministic computing


Updating rule

- one – dimensional chain, x : discrete lattice sites
- at each x : red and green right movers and left movers (4 different species at each site)
- at each time step: configuration for right(left) movers moves one position to the right(left)
- if two single particles meet: colors are exchanged





Updating rule


- at each time step configuration for right(left) movers moves one position to the right(left), **periodicity in x**
- if **precisely two single particles** meet at a site : colors are exchanged



Half filled ground states




Soliton



black line : two right movers with different colors

Soliton separates different vacua



Probabilistic cellular automaton

Probability distribution for initial configurations

(or other probabilistic boundary condition)

Probabilistic formalism for cellular automata

Generalized Ising model:

$$w[s] = Z^{-1} \exp(-S[s]) b(s_{in}, s_f)$$

$$S = -\frac{\beta}{2} \sum_{x,t} s(t,x) \left[s(t+1, x+1) + \sigma s(t+1, x-1) \right]$$

limit : beta to infinity , sigma to zero :
only one possibility for change , **unique jump**

probabilistic
aspects only in
boundary term :

$$b(s_{in}, s_f) = \bar{f}_f(s_f) f_{in}(s_{in})$$

Functional integral for cellular automata

Generalized Ising model:

$$w[s] = Z^{-1} \exp(-S[s]) b(s_{in}, s_f)$$

$$S = -\frac{\beta}{2} \sum_{x,t} s(t,x) \left[s(t+1, x+1) + \sigma s(t+1, x-1) \right]$$

limit : beta to infinity , sigma to zero :

only one possibility for change , **unique jump**

Functional renormalization for cellular automata

Probabilistic computing with static memory materials ?

- Let general equilibrium classical statistics transport information from one layer to the next
- Simulation, with D. Sexty

Static memory materials

Generalized Ising model:

$$w[s] = Z^{-1} \exp(-S[s]) b(s_{in}, s_f)$$


$$S = -\frac{\beta}{2} \sum_{x,t} s(t, x) \left[s(t+1, x+1) + \sigma s(t+1, x-1) \right]$$

Boundary term :


$$b(s_{in}, s_f) = \bar{f}_f(s_f) f_{in}(s_{in})$$

Classical interference

Depending on boundary conditions :




Positive
interference



Negative
interference

*Static memory material for
two dimensional Ising spins
on Euclidean square lattice
can describe propagation of Weyl fermion
in two- dimensional Minkowski space*

*Probabilistic cellular automaton is
equivalent to quantum field theory
for fermions with interactions*



Discrete fermion model in 1+1 dimensions

Grassmann functional integral

$$Z = \int \mathcal{D}\psi \exp(-S[\psi]) = \int \mathcal{D}\psi w[\psi], \quad S = \sum_t \mathcal{L}(t)$$

$$\mathcal{L}(t) = - \sum_x \left\{ \bar{\psi}_{R\alpha}(t + \varepsilon, x + \varepsilon) \psi_{R\alpha}(t, x) + \bar{\psi}_{L\alpha}(t + \varepsilon, x - \varepsilon) \psi_{L\alpha}(t, x) \right. \\ \left. - \left[\bar{\psi}_{R\alpha}(t, x) \psi_{R\alpha}(t, x) + \bar{\psi}_{L\alpha}(t, x) \psi_{L\alpha}(t, x) + \bar{D}(x) \right] (1 + \bar{D}(x)) \right\}$$

$$\bar{D} = -(\bar{\psi}_{R1} \bar{\psi}_{L2} - \bar{\psi}_{R2} \bar{\psi}_{L1})(\psi_{R1} \psi_{L2} - \psi_{R2} \psi_{L1}) - (\bar{\psi}_{R1} \bar{\psi}_{L1} + \bar{\psi}_{R2} \bar{\psi}_{L2})(\psi_{R1} \psi_{L1} + \psi_{R2} \psi_{L2})$$

General bit- fermion map

- Fermion model in terms of Grassmann functional integral is equivalent to generalized Ising model
- Isomorphism for evolution and observables
- Map is general. But not always positive weight distribution for Ising type model, and unitary evolution.

Continuum limit

$$S = \int_{t,x} \{ \bar{\psi}_{R\alpha}(t,x)(\partial_t + \partial_x)\psi_{R\alpha}(t,x) + \bar{\psi}_{L\alpha}(t,x)(\partial_t - \partial_x)\psi_{L\alpha}(t,x) + 2\bar{D}(t,x) \}$$

$$\bar{D} = -(\bar{\psi}_{R1}\bar{\psi}_{L2} - \bar{\psi}_{R2}\bar{\psi}_{L1})(\psi_{R1}\psi_{L2} - \psi_{R2}\psi_{L1}) - (\bar{\psi}_{R1}\bar{\psi}_{L1} + \bar{\psi}_{R2}\bar{\psi}_{L2})(\psi_{R1}\psi_{L1} + \psi_{R2}\psi_{L2})$$

$$\begin{aligned}(\partial_t + \partial_x)\psi(t,x) &= \frac{1}{\varepsilon} [\psi(t,x) - \psi(t-\varepsilon, x-\varepsilon)] \\ (\partial_t - \partial_x)\psi(t,x) &= \frac{1}{\varepsilon} [\psi(t,x) - \psi(t-\varepsilon, x+\varepsilon)]\end{aligned}$$

$$\int dt \int dx = \int_{t,x} = 2\varepsilon^2 \sum_{t,x}$$

$$\psi(t,x) = \sqrt{2\varepsilon} \psi_N(t,x)$$

Lorentz symmetry

Dirac spinor

$$\psi_a = \begin{pmatrix} \psi_{Ra} \\ \psi_{La} \end{pmatrix}, \quad \bar{\psi}_a = (\bar{\psi}_{La}, -\bar{\psi}_{Ra})$$

Action

$$S = - \int_{t,x} \left\{ \bar{\psi}_a \gamma^\mu \partial_\mu \psi_a + \frac{1}{2} \bar{\psi}_a \gamma^\mu \psi_a \bar{\psi}_b \gamma_\mu \psi_b + \frac{1}{2} \bar{\psi}_a \gamma^\mu \psi_b \epsilon^{ab} \bar{\psi}_c \gamma_\mu \psi_d \epsilon^{cd} \right\}$$

$$\gamma^0 = -i\tau_2, \quad \gamma_1 = \tau_1, \quad \{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu}$$

Infinitesimal Lorentz transformation

$$\delta\psi = -\eta \Sigma^{01} \psi, \quad \delta\bar{\psi} = \eta \bar{\psi} \Sigma^{01}$$

$$\Sigma^{01} = \frac{1}{4} [\gamma^0, \gamma^1] = \frac{1}{2} \tau_3$$

Quantum field theory for fermions with interactions

- Thirring model with two colors
- Particular value of coupling

$$S = - \int_{t,x} \left\{ \bar{\psi}_a \gamma^\mu \partial_\mu \psi_a + \frac{1}{2} \bar{\psi}_a \gamma^\mu \psi_a \bar{\psi}_b \gamma_\mu \psi_b + \frac{1}{2} \bar{\psi}_a \gamma^\mu \psi_b \epsilon^{ab} \bar{\psi}_c \gamma_\mu \psi_d \epsilon^{cd} \right\}$$

The background of the slide is a dark, almost black, field with a series of bright, white, and grey light rays or beams emanating from a point on the left side, creating a sense of depth and movement. The rays are slightly blurred, giving the impression of light traveling through a medium.

Quantum mechanics

from classical statistics

Quantum mechanics from classical statistics

For **particular** quantum model:

Isomorphism between classical statistics

(probabilistic cellular automaton,
generalized Ising model)

and quantum mechanics

(many body quantum system for fermions)

Equivalence

- Expectation values of all observables are the same in both models
- Two equivalent descriptions of the same physical reality

Can quantum physics be described by classical probabilities ?

“No go” theorems

Bell , Clauser , Horne , Shimony , Holt

implicit assumption : use of classical correlation function for all correlations between measurements

Kochen , Specker

assumption : unique map from quantum operators to classical observables

Probabilistic cellular automaton

Probabilistic initial condition: Specify

at initial time t_{in} for each bit configuration $\bar{\rho}$
a probability $p_{\bar{\rho}}(t_{\text{in}})$

Evolution: every given configuration $\bar{\rho}$ at t_{in}
propagates at t to a configuration $\tau(t, \bar{\rho})$



$$p_{\tau}(t) = p_{\bar{\rho}(\tau)}(t_{\text{in}})$$

Updating rule: specifies

$$\tau(t + \varepsilon, \rho(t))$$

Wave function for probabilistic cellular automaton

Probability distribution: at every time t a bit configuration τ occurs with probability $p_{\tau}(t)$

Real wave function $q(t)$: probability amplitude

$$p_{\tau}(t) = (q_{\tau}(t))^2$$

$$q_{\tau}(t)q_{\tau}(t) = 1$$

N – component unit vector

Deterministic and probabilistic cellular automaton

- Deterministic CA : sharp wave function

$$q_{\rho}(t_{\text{in}}) = \delta_{\rho, \bar{\rho}}$$

- Probabilistic CA : arbitrary wave function

Particle wave duality

Particle aspect:

- Bits: yes/no decisions
- Possible measurement values 1 or 0

Discrete spectrum of observables

Wave aspect : continuous wave function

more generally: continuity of probabilistic information

Step evolution operator

- Evolution for basic time step is encoded in the step evolution operator

$$q(t + \varepsilon) = \hat{S}(t)q(t) \quad q_\tau(t + \varepsilon) = \hat{S}_{\tau\rho}(t)q_\rho(t)$$

- Contains the updating rule for CA

$$\hat{S}_{\tau\rho}(t) = \delta_{\tau, \overline{\rho}(\rho)} = \delta_{\overline{\rho}(\tau), \rho}$$

$$q_\tau(t + \varepsilon) = q_{\overline{\rho}(\tau)}(t), \quad p_\tau(t + \varepsilon) = p_{\overline{\rho}(\tau)}(t)$$

Step evolution operator

- Sequence of kinetic (free) and interaction part

$$\hat{S} = \hat{S}_{\text{int}} \hat{S}_{\text{free}}$$

- Local interaction

$$\hat{S}_{\text{int}} = \hat{S}_i(x_{\text{in}}) \otimes \hat{S}_i(x_{\text{in}} + \varepsilon) \otimes \hat{S}_i(x_{\text{in}} + 2\varepsilon) \otimes \dots$$

- (1) at each time step configuration for right(left) movers moves one position to the right(left),
- (2) if **precisely two single particles** meet at a site : colors are exchanged

Annihilation and creation operators

Step evolution operator can be written in terms of fermionic annihilation and creation operators

$$\{a_{\gamma}^{\dagger}(x), a_{\delta}(y)\} = \delta_{\gamma\delta}\delta_{xy} \quad \{a_{\gamma}(x), a_{\delta}(y)\} = \{a_{\gamma}^{\dagger}(x), a_{\delta}^{\dagger}(y)\} = 0$$

$$\hat{S}_i(x) = \exp \left\{ \frac{i\pi}{2} [a_{R1}^{\dagger}(x)a_{R2}(x) - a_{R2}^{\dagger}(x)a_{R1}(x)] [a_{L1}^{\dagger}(x)a_{L2}(x) - a_{L2}^{\dagger}(x)a_{L1}(x)] \right\}$$

$$\hat{S}_{\text{free}} = \hat{S}_1^{(R)} \otimes \hat{S}_2^{(R)} \otimes \hat{S}_1^{(L)} \otimes \hat{S}_2^{(L)}$$

$$\hat{S}_a^{(R,L)} = N \left[\exp \left\{ \sum_x a^{\dagger}(x \pm \varepsilon) [a(x) - a(x \pm \varepsilon)] \right\} \right]$$

Hamiltonian

- Define H by $\hat{S} = \exp(-i\varepsilon H)$

- Interpolating continuous time evolution

$$q(t_2) = U(t_2, t_1)q(t_1) \quad U(t_1, t_2) = \exp(-i(t_1 - t_2)H)$$

- Agrees with discrete evolution for $t_{\text{in}} + m\varepsilon$

- Schrödinger equation $i\partial_t q = Hq$

Continuum limit

- Hamiltonian simplifies in the continuum limit

$$H = H_{\text{free}} + H_{\text{int}} + \Delta H$$

$$\Delta H = \mathcal{O}(\varepsilon[H_{\text{int}}, H_{\text{free}}])$$

- Standard form of Hamiltonian for fermions

$$H_{\text{free}} = \frac{i}{\varepsilon} \int dx \sum_a \left\{ a_{La}^\dagger(x) \partial_x a_{La}(x) - a_{Ra}^\dagger(x) \partial_x a_{Ra}(x) \right\}$$

$$H_{\text{int}} = - \frac{\pi}{2\varepsilon^2} \int dx [a_{R1}^\dagger a_{R2} - a_{R2}^\dagger a_{R1}] [a_{L1}^\dagger a_{L2} - a_{L2}^\dagger a_{L1}]$$

General bit fermion map

- Isomorphism between generalized Ising model and Grassmann functional integral
- Based on identical step evolution operator for both models, with associated map of observables
- In our case: proof that discrete Thirring model with two colors has the same step evolution operator as the cellular automaton

$$\mathcal{L}(t) = - \sum_x \left\{ \bar{\psi}_{R\alpha}(t + \varepsilon, x + \varepsilon) \psi_{R\alpha}(t, x) + \bar{\psi}_{L\alpha}(t + \varepsilon, x - \varepsilon) \psi_{L\alpha}(t, x) \right. \\ \left. - \left[\bar{\psi}_{R\alpha}(t, x) \psi_{R\alpha}(t, x) + \bar{\psi}_{L\alpha}(t, x) \psi_{L\alpha}(t, x) + \overline{D}(x) \right] (1 + \overline{D}(x)) \right\}$$

Quantum formalism for classical statistics

- Formalism for information transport from one hypersurface to the next:
- Classical wave functions and density matrix
- Transfer matrix formalism : Heisenberg picture
- Wave functions : Schrödinger picture
- Non commuting operators for observables
- Quantum rules from classical statistical rules

Conclusion

- Particular quantum field theory for interacting fermions is equivalent to the classical statistical model of a particular probabilistic cellular automaton.
- Example for quantum mechanics from classical statistics
- Useful for simulating fermionic models and understanding of statistical properties of cellular automata?

end

Reduction of wave function


- Reduction of wave function is a convenient technical method to describe conditional probabilities
- This must not be a physical process during the measurement

conditional probability

sequences of events(measurements)
are described by
conditional probabilities

*both in classical statistics
and in quantum statistics*

$w(t_1)$



:

not very suitable
for statement , if here and now
a pointer falls down

Schrödinger's cat



conditional probability :
if nucleus decays
then cat dead with $w_c = 1$
(reduction of wave function)

structural elements of quantum mechanics

unitary time evolution



h

Simple conversion factor for units

i

presence of complex structure

$$[A, B] = C$$

non – commuting operators
are necessary to represent
observables in
incomplete statistics

correlation and operator product

- Classical statistical systems admit many product structures of observables
- Many different definitions of correlation functions possible , not only classical correlation !
- Type of measurement determines correct selection of correlation function !
- Ideal quantum measurement should be compatible with information in quantum subsystem

Deterministic evolution – probabilistic interpretation

- quantum mechanics arises from

quantum subsystems

- subsystems are genuinely probabilistic
- part of information is lost by focus on subsystem
- partially "integrating out" degrees of freedom

Determinism vs. Probabilism



“ Does god throw dices ? ”

... an old dispute

Gott würfelt



Gott würfelt nicht



“Es scheint hart, dem Herrgott in die Karten zu gucken. Aber dass er würfelt und sich telepatischer Mittel bedient (wie es ihm von der gegenwärtigen Quantentheorie zugemutet wird), kann ich keinen Augenblick glauben..”

Einstein: Brief an Cornelius Lanczos am 21. März 1942

not today's topic

Gott würfelt

Gott würfelt nicht



humans can only deal with probabilities



determinism vs. probabilism

my personal view :

- determinism not needed
- one can start with probabilistic theory and probabilistic evolution
- nevertheless : deterministic evolution is a possible option