

*A quantum information perspective on relativistic fluid
dynamics and quantum fields out-of-equilibrium*

Stefan Floerchinger (Heidelberg U.)

Nikhef,
Amsterdam, 12 December 2019.



UNIVERSITÄT
HEIDELBERG
ZUKUNFT
SEIT 1386



STRUCTURES
CLUSTER OF EXCELLENCE

Fluid dynamics




- long distances, long times or strong enough interactions
- quantum fields form a fluid!
- needs **macroscopic** fluid properties
 - equation of state $p(T, \mu)$
 - shear viscosity $\eta(T, \mu)$
 - bulk viscosity $\zeta(T, \mu)$
 - relaxation times, ...
- *ab initio* calculation of transport properties difficult but in principle fixed by **microscopic** properties encoded in lagrangian
- standard model of high energy nuclear collisions based on relativistic dissipative fluid dynamics
- ongoing experimental and theoretical effort to understand this better

Fluid description of high energy nuclear collisions

[Devetak, Dubla, Floerchinger, Grossi, Masciocchi, Mazeliauskas & Selyuzhenkov, 1909.10485]

Identified particle transverse momentum spectra at the LHC




Entropy and information

[Claude Shannon (1948)]


- consider a random variable x with probability distribution $p(x)$
- information content or “surprise” associated with outcome x

$$i(x) = -\ln p(x)$$




- entropy is expectation value of information content


$$S = \langle i(x) \rangle = -\sum_x p(x) \ln p(x)$$



$$S = 0$$



$$S = \ln(2)$$



$$S = 2 \ln(2)$$

Entropy at thermal equilibrium


- micro canonical ensemble: **maximal entropy** S for given **conserved quantities** E, N in given volume V
- **universality** at equilibrium
- starting point for development of thermodynamics ...

$$S(E, N, V), \quad dS = \frac{1}{T}dE - \frac{\mu}{T}dN + \frac{p}{T}dV$$

- ... grand canonical ensemble with density operator ...

$$\rho = \frac{1}{Z} e^{-\frac{1}{T}(H - \mu N)}$$

- ... Matsubara formalism for quantum fields ...



Ideal fluid dynamics

- thermal equilibrium

$$T^{\mu\nu} = \epsilon u^\mu u^\nu + p(u^\mu u^\nu + g^{\mu\nu}), \quad N^\mu = n u^\mu, \quad s^\mu = s u^\mu$$

- fluid velocity u^μ
- thermodynamic equation of state $p(T, \mu)$ with $dp = sdT + nd\mu$
- local thermal equilibrium approximation: $u^\mu(x), T(x), \mu(x)$
- neglect gradients: lowest order of a derivative expansion
- evolution of $u^\mu(x), T(x)$ and $\mu(x)$ from conservation laws

$$\nabla_\mu T^{\mu\nu}(x) = 0, \quad \nabla_\mu N^\mu(x) = 0.$$

- entropy current also conserved

$$\nabla_\mu s^\mu(x) = 0.$$

Out-of-equilibrium

- is non-equilibrium dynamics also governed by information?
- approach to equilibrium
- universality

Entropy in quantum theory

[John von Neumann (1932)]

$$S = -\text{Tr}\{\rho \ln \rho\}$$

- based on the quantum density operator ρ
- for pure states $\rho = |\psi\rangle\langle\psi|$ one has $S = 0$
- for mixed states $\rho = \sum_j p_j |j\rangle\langle j|$ one has $S = -\sum_j p_j \ln p_j > 0$
- unitary time evolution conserves entropy

$$-\text{Tr}\{(U\rho U^\dagger) \ln(U\rho U^\dagger)\} = -\text{Tr}\{\rho \ln \rho\} \quad \rightarrow \quad S = \text{const.}$$

- quantum information is globally conserved

Dissipative relativistic fluid dynamics

- approximate description of quantum field dynamics
- local dissipation = local entropy production

$$\nabla_{\mu} s^{\mu}(x) > 0$$

- e. g. in Navier-Stokes approximation

$$\nabla_{\mu} s^{\mu} = \frac{1}{T} [2\eta\sigma_{\mu\nu}\sigma^{\mu\nu} + \zeta(\nabla_{\rho} u^{\rho})^2] \geq 0$$

- crucial difference to quantum field theory: entropy not conserved

What is an entropy current?

- *can not* be density of global von-Neumann entropy for closed system

$$\int_{\Sigma} d\Sigma_{\mu} s^{\mu}(x) \neq -\text{Tr} \{ \rho \ln \rho \}$$


- kinetic theory for weakly coupled (quasi-) particles [Boltzmann (1890)]

$$s^{\mu}(x) = - \int \frac{d^3p}{p^0} \{ p^{\mu} f(x, p) \ln f(x, p) \}$$

- molecular chaos: keep only single particle distribution $f(x, p)$
- how to go beyond weak coupling / quasiparticles?
- aim: local notion of entropy in QFT

Entropy and entanglement

- consider a split of a quantum system into two $A + B$



- reduced density operator for system A

$$\rho_A = \text{Tr}_B\{\rho\}$$

- entropy associated with subsystem A

$$S_A = -\text{Tr}_A\{\rho_A \ln \rho_A\}$$


- pure **product** state $\rho = \rho_A \otimes \rho_B$ leads to $S_A = 0$
- pure **entangled** state $\rho \neq \rho_A \otimes \rho_B$ leads to $S_A > 0$
- S_A is called **entanglement entropy**

Why is entanglement interesting?

- Can quantum-mechanical description of physical reality be considered complete? [Einstein, Podolsky, Rosen (1935), Bohm (1951)]

$$\begin{aligned} |\psi\rangle &= \frac{1}{\sqrt{2}} (|\uparrow\rangle_A |\downarrow\rangle_B - |\downarrow\rangle_A |\uparrow\rangle_B) \\ &= \frac{1}{\sqrt{2}} (|\rightarrow\rangle_A |\leftarrow\rangle_B - |\leftarrow\rangle_A |\rightarrow\rangle_B) \end{aligned}$$

- Bertlemann's socks and the nature of reality [Bell (1980)]



Bell's inequalities and Bell tests

[John Stewart Bell (1966)]

- most popular version [Clauser, Horne, Shimony, Holt (1969)]

$$S = |E(a, b) - E(a, b') + E(a', b) + E(a', b')| \leq 2$$

holds for local hidden variable theories

- expectation value of product of two observables

$$E(a, b) = \langle A(a)B(b) \rangle$$

with possible values $A = \pm 1$, $B = \pm 1$.

- depending on measurement settings a , a' and b , b' respectively
- quantum mechanical bound is $S \leq 2\sqrt{2}$
- experimental values $2 < S \leq 2\sqrt{2}$ rule out local hidden variables
- one measurement setting but at different times [Leggett, Garg (1985)]

Entanglement in high energy (QCD) physics

[. . . , Elze (1996), Kovner, Lublinsky (2015), Kharzeev & Levin (2017), Berges, Floerchinger & Venugopalan (2017), Shuryak & Zahed (2017), Kovner, Lublinsky, Serino (2018), Baker & Kharzeev (2018), Tu, Kharzeev & Ullrich (2019), Armesto, Dominguez, Kovner, Lublinsky, Skokov (2019), . . .]

- entanglement of *quantum fields* instead of *particles*
- entanglement on sub-nucleonic scales
- entanglement in non-Abelian gauge theory / color / confinement
- discussions in mathematical physics [e. g. Witten (2018)]
- connections to black holes and holography [Ryu & Takayanagi (2006)]
- thermalization in closed quantum systems

Classical statistics

- consider system of two random variables x and y
- joint probability $p(x, y)$, joint entropy

$$S = - \sum_{x,y} p(x, y) \ln p(x, y)$$

- reduced or marginal probability $p(x) = \sum_y p(x, y)$
- reduced or marginal entropy

$$S_x = - \sum_x p(x) \ln p(x)$$

- one can prove: **joint entropy is greater than** or equal to **reduced entropy**

$$S \geq S_x$$

- **globally pure** state $S = 0$ is also **locally pure** $S_x = 0$

Quantum statistics

- consider system with two subsystems A and B
- combined state ρ , combined or full entropy

$$S = -\text{Tr}\{\rho \ln \rho\}$$

- reduced density matrix $\rho_A = \text{Tr}_B\{\rho\}$
- reduced or entanglement entropy

$$S_A = -\text{Tr}_A\{\rho_A \ln \rho_A\}$$

- for quantum systems **entanglement makes a difference**

$$S \not\equiv S_A$$

- **coherent information** $I_{B \rangle A} = S_A - S$ can be **positive!**
- **globally pure** state $S = 0$ can be **locally mixed** $S_A > 0$

Entanglement entropy in quantum field theory

- entanglement entropy of region A is a local notion of entropy

$$S_A = -\text{tr}_A \{ \rho_A \ln \rho_A \} \quad \rho_A = \text{tr}_B \{ \rho \}$$

- however, it is infinite already in vacuum state

$$S_A = \frac{\text{const}}{\epsilon^{d-2}} \int_{\partial A} d^{d-2} \sigma \sqrt{h} + \text{subleading divergences} + \text{finite}$$

- UV divergence proportional to entangling surface
- quantum fields are very strongly entangled already in vacuum
- Theorem [Reeh & Schlieder (1961)]: local operators in region A can create all particle states

Relative entropy

- **relative entropy** of two density matrices

$$S(\rho|\sigma) = \text{tr} \{ \rho (\ln \rho - \ln \sigma) \}$$

- measures how well state ρ can be distinguished from a model σ
- Gibbs inequality: $S(\rho|\sigma) \geq 0$
- $S(\rho|\sigma) = 0$ if and only if $\rho = \sigma$
- quantum generalization of Kullback-Leibler divergence

Relative entanglement entropy

- consider now reduced density matrices

$$\rho_A = \text{Tr}_B\{\rho\}, \quad \sigma_A = \text{Tr}_B\{\sigma\}$$

- define **relative entanglement entropy**

$$S_A(\rho|\sigma) = \text{Tr}\{\rho_A (\ln \rho_A - \ln \sigma_A)\}$$

- measures how well ρ is represented by σ locally in region A
- UV divergences cancel: contains real physics information
- well defined in quantum field theory [Araki (1977)]
[see also works by Casini, Myers, Lashkari, Witten, Liu, ...]

An approximate local description

- consider non-equilibrium situation with
 - true density matrix ρ
 - local equilibrium approximation

$$\sigma = \frac{1}{Z} e^{-\int d\Sigma_\mu \{\beta_\nu(x) T^{\mu\nu} + \alpha(x) N^\mu\}}$$

- reduced density matrices $\rho_A = \text{Tr}_B\{\rho\}$ and $\sigma_A = \text{Tr}_B\{\sigma\}$
- σ is very good model for ρ in region A when

$$S_A = \text{Tr}_A\{\rho_A(\ln \rho_A - \ln \sigma_A)\} \rightarrow 0$$

- does *not* imply that globally $\rho = \sigma$

Monotonicity of relative entropy

- monotonicity of relative entropy

$$S(\mathcal{N}(\rho)|\mathcal{N}(\sigma)) \leq S(\rho|\sigma)$$

with \mathcal{N} completely positive, trace-preserving map

- \mathcal{N} unitary evolution

$$S(\mathcal{N}(\rho)|\mathcal{N}(\sigma)) = S(\rho|\sigma)$$

- \mathcal{N} open system evolution with generation of entanglement to environment

$$S(\mathcal{N}(\rho)|\mathcal{N}(\sigma)) < S(\rho|\sigma)$$

Local form of second law

- for small volume $A \rightarrow 0$ (hypothesis)

$$S_A(\rho|\sigma) = \int_A d\Sigma_\mu s^\mu(\rho|\sigma)$$

- local form of second law of thermodynamics

$$\nabla_\mu s^\mu(\rho|\sigma) \leq 0$$


- **relative entanglement entropy between ρ and any state, in particular thermal state σ is non-increasing**

Quantum field dynamics

- new hypothesis


local dissipation = *quantum entanglement generation*

- quantum information is spread
- locally, quantum state approaches mixed state form
- full loss of *local* quantum information = *local* thermalization



Local equilibrium & partition function

[Floerchinger, JHEP 1609, 099 (2016)]



- local equilibrium with $T(x)$ and $u^\mu(x)$

$$\beta^\mu(x) = \frac{u^\mu(x)}{T(x)}$$

- represent partition function as functional integral with periodicity

$$\phi(x^\mu - i\beta^\mu(x)) = \pm\phi(x^\mu)$$

- partition function $Z[J]$, Schwinger functional $W[J]$ in Euclidean

$$Z[J] = e^{W_E[J]} = \int D\phi e^{-S_E[\phi] + \int_x J\phi}$$

One-particle irreducible or quantum effective action

- in Euclidean domain $\Gamma[\phi]$ defined by Legendre transform

$$\Gamma_E[\Phi] = \int_x J_a(x) \Phi_a(x) - W_E[J]$$

with expectation values

$$\Phi_a(x) = \frac{1}{\sqrt{g(x)}} \frac{\delta}{\delta J_a(x)} W_E[J]$$

- **Euclidean** field equation

$$\frac{\delta}{\delta \Phi_a(x)} \Gamma_E[\Phi] = \sqrt{g(x)} J_a(x)$$

resembles classical equation of motion for $J = 0$

- need **analytic continuation** to obtain a viable equation of motion

Entropy production

[Floerchinger, JHEP 1609, 099 (2016)]

- variational principle with effective dissipation from analytic continuation
- analysis of general covariance leads to entropy current and local entropy production


$$\nabla_{\mu} s^{\mu} = \frac{1}{\sqrt{g}} \frac{\delta \Gamma_D}{\delta \Phi_a} \Big|_{\text{ret}} \beta^{\lambda} \partial_{\lambda} \Phi_a + \beta_{\mu} \nabla_{\nu} \left(-\frac{2}{\sqrt{g}} \frac{\delta \Gamma_D}{\delta g_{\mu\nu}} \Big|_{\text{ret}} \right)$$

- can likely be understood as entanglement generation

Thermalization beyond collisions

- quantum fields can be locally thermal without collisions
- horizons: black holes, de-Sitter space
- space-time dynamics of entanglement

Entanglement, QCD strings and thermalization



- hadronization in Lund string model (e. g. PYTHIA)
- reduced density matrix for region A

$$\rho_A = \text{Tr}_B\{\rho\}$$


- **entanglement entropy**

$$S_A = -\text{Tr}_A\{\rho_A \ln \rho_A\}$$

- could this lead to thermal-like effects?

The thermal model puzzle

- elementary particle collision experiments such as $e^+ e^-$ collisions show some thermal-like features
- particle multiplicities well described by thermal model



[Becattini, Casterina, Milov & Satz, EPJC 66, 377 (2010)]

- conventional thermalization by collisions unlikely
- more thermal-like features difficult to understand in PYTHIA
[Fischer, Sjöstrand (2017)]
- alternative explanations needed

Microscopic model

- QCD in 1+1 dimensions described by 't Hooft model

$$\mathcal{L} = -\bar{\psi}_i \gamma^\mu (\partial_\mu - ig\mathbf{A}_\mu) \psi_i - m_i \bar{\psi}_i \psi_i - \frac{1}{2} \text{tr} \mathbf{F}_{\mu\nu} \mathbf{F}^{\mu\nu}$$

- fermionic fields ψ_i with sums over flavor species $i = 1, \dots, N_f$
- $SU(N_c)$ gauge fields \mathbf{A}_μ with field strength tensor $\mathbf{F}_{\mu\nu}$
- gluons are not dynamical in two dimensions
- gauge coupling g has dimension of mass
- non-trivial, interacting theory, cannot be solved exactly
- spectrum of excitations known for $N_c \rightarrow \infty$ with $g^2 N_c$ fixed
[t Hooft (1974)]

Schwinger model

- QED in 1+1 dimension


$$\mathcal{L} = -\bar{\psi}_i \gamma^\mu (\partial_\mu - iqA_\mu) \psi_i - m_i \bar{\psi}_i \psi_i - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}$$

- geometric confinement
- U(1) charge related to string tension $q = \sqrt{2\sigma}$
- for single fermion one can **bosonize theory** exactly
[Coleman, Jackiw, Susskind (1975)]

$$S = \int d^2x \sqrt{g} \left\{ -\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} M^2 \phi^2 - \frac{m q e^\gamma}{2\pi^{3/2}} \cos(2\sqrt{\pi}\phi + \theta) \right\}$$

- Schwinger bosons are dipoles $\phi \sim \bar{\psi}\psi$
- scalar mass related to U(1) charge by $M = q/\sqrt{\pi} = \sqrt{2\sigma/\pi}$
- massless Schwinger model $m = 0$ leads to free bosonic theory

Expanding string solution 1



- external quark-anti-quark pair on trajectories $z = \pm t$
- coordinates: Bjorken time $\tau = \sqrt{t^2 - z^2}$, rapidity $\eta = \text{arctanh}(z/t)$
- metric $ds^2 = -d\tau^2 + \tau^2 d\eta^2$
- symmetry with respect to longitudinal boosts $\eta \rightarrow \eta + \Delta\eta$

Expanding string solution 2

- Schwinger boson field depends only on τ

$$\bar{\phi} = \bar{\phi}(\tau)$$

- equation of motion

$$\partial_\tau^2 \bar{\phi} + \frac{1}{\tau} \partial_\tau \bar{\phi} + M^2 \bar{\phi} = 0.$$

- Gauss law: electric field $E = q\phi/\sqrt{\pi}$ must approach the U(1) charge of the external quarks $E \rightarrow q_e$ for $\tau \rightarrow 0_+$

$$\bar{\phi}(\tau) \rightarrow \frac{\sqrt{\pi}q_e}{q} \quad (\tau \rightarrow 0_+)$$

- solution of equation of motion [Loshaj, Kharzeev (2011)]

$$\bar{\phi}(\tau) = \frac{\sqrt{\pi}q_e}{q} J_0(M\tau)$$

Gaussian states

- theories with quadratic action often have Gaussian density matrix
- fully characterized by field expectation values

$$\bar{\phi}(x) = \langle \phi(x) \rangle, \quad \bar{\pi}(x) = \langle \pi(x) \rangle$$

and connected two-point correlation functions, e. g.

$$\langle \phi(x)\phi(y) \rangle_c = \langle \phi(x)\phi(y) \rangle - \bar{\phi}(x)\bar{\phi}(y)$$

- if ρ is Gaussian, also reduced density matrix ρ_A is Gaussian

Entanglement entropy for Gaussian state

- entanglement entropy of Gaussian state in region A
[Berges, Floerchinger, Venugopalan, JHEP 1804 (2018) 145]

$$S_A = \frac{1}{2} \text{Tr}_A \{ D \ln(D^2) \}$$


- operator trace over region A only
- matrix of correlation functions

$$D(x, y) = \begin{pmatrix} -i\langle \phi(x)\pi(y) \rangle_c & i\langle \phi(x)\phi(y) \rangle_c \\ -i\langle \pi(x)\pi(y) \rangle_c & i\langle \pi(x)\phi(y) \rangle_c \end{pmatrix}$$

- involves connected correlation functions of field $\phi(x)$ and canonically conjugate momentum field $\pi(x)$
- expectation value $\bar{\phi}$ does not appear explicitly
- coherent states and vacuum have equal entanglement entropy S_A

Local density matrix and temperature in expanding string

[Berges, Floerchinger, Venugopalan, *Thermal excitation spectrum from entanglement in an expanding quantum string*, PLB778, 442 (2018)]



- Bjorken time $\tau = \sqrt{t^2 - z^2}$, rapidity $\eta = \text{arctanh}(z/t)$
- **local density matrix thermal at early times as result of entanglement**

$$T(\tau) = \frac{\hbar}{2\pi\tau}$$

- Hawking-Unruh temperature in Rindler space $T(x) = \frac{\hbar c}{2\pi x}$

Physics picture

- coherent state at early time contains entangled pairs of quasi-particles with opposite wave numbers
- on finite rapidity interval $(-\Delta\eta/2, \Delta\eta/2)$ in- and out-flux of quasi-particles with thermal distribution via boundaries
- technically limits $\Delta\eta \rightarrow \infty$ and $M\tau \rightarrow 0$ do not commute
 - $\Delta\eta \rightarrow \infty$ for any finite $M\tau$ gives pure state
 - $M\tau \rightarrow 0$ for any finite $\Delta\eta$ gives thermal state with $T = 1/(2\pi\tau)$

Testing the picture

- explicit calculations in non-equilibrium QFT
- experimental tests with high-energy collisions
- explicit calculations in holography
- explicit calculations in small dimensions with tensor networks
- quantum simulations with universal quantum computers
- quantum simulation with ultracold atoms

Entropic uncertainty relations

Heisenberg / Robertson uncertainty relation [Robertson (1929)]

$$\sigma(X)\sigma(Z) \geq \frac{1}{2} |\langle \psi | [X, Z] | \psi \rangle|$$

Entropic uncertainty relations [Maassen & Uffink (1988), Frank & Lieb (2012)]

$$H(X) + H(Z) \geq \ln \frac{1}{c} + S(\rho)$$

- Shannon information entropy for measurement outcome

$$H(X) = - \sum_x p(x) \ln p(x)$$

- von-Neumann entropy

$$S(\rho) = -\text{Tr}\{\rho \ln \rho\}$$

- maximal overlap between basis states

$$c = \max_{x,z} |\langle x|z \rangle|^2$$

Entanglement and entropic uncertainty relations

[Berta et al. (2010)]

- side information from entanglement with system B

$$H(X_A|X_B) + H(Z_A|Z_B) \geq \ln \frac{1}{c} + S(A|B)$$

- use measurement on B to infer outcome on A
- quantum conditional entropy can be negative for positive coherent information

$$S(A|B) = S(\rho) - S(\rho_B) = -I_{A>B}$$


- experiments with cold atoms [with M. Gärttner and M. Oberthaler]
- towards test of *local dissipation = quantum entanglement generation*
- towards test of entanglement in horizon physics
- more applications in nuclear and high energy physics to be explored

Conclusions

- new perspectives on relativistic fluids from quantum information theory
- relative entanglement entropy useful to describe local thermalization
- quantum field theoretic description of relativistic fluid dynamics with two density matrices
 - true density matrix ρ evolves unitary
 - fluid model σ agrees locally but evolves non-unitary
- local thermalization without collisions possible
- excitations in expanding QCD strings locally thermal at early times

Backup


Rapidity interval



- consider rapidity interval $(-\Delta\eta/2, \Delta\eta/2)$ at fixed Bjorken time τ
- entanglement entropy does not change by unitary time evolution with endpoints kept fixed
- can be evaluated equivalently in interval $\Delta z = 2\tau \sinh(\Delta\eta/2)$ at fixed time $t = \tau \cosh(\Delta\eta/2)$
- need to solve eigenvalue problem with correct **boundary conditions**

Bosonized massless Schwinger model

- entanglement entropy understood numerically for free massive scalars [Casini, Huerta (2009)]
- entanglement entropy density $dS/d\Delta\eta$ for bosonized massless Schwinger model ($M = \frac{q}{\sqrt{\pi}}$)



Conformal limit

- For $M\tau \rightarrow 0$ one has conformal field theory limit

[Holzhey, Larsen, Wilczek (1994)]

$$S(\Delta z) = \frac{c}{3} \ln(\Delta z/\epsilon) + \text{constant}$$

with small length ϵ acting as UV cutoff.

- Here this implies

$$S(\tau, \Delta\eta) = \frac{c}{3} \ln(2\tau \sinh(\Delta\eta/2)/\epsilon) + \text{constant}$$

- Conformal charge $c = 1$ for free massless scalars or Dirac fermions.
- Additive constant not universal but entropy density is

$$\begin{aligned} \frac{\partial}{\partial \Delta\eta} S(\tau, \Delta\eta) &= \frac{c}{6} \coth(\Delta\eta/2) \\ &\rightarrow \frac{c}{6} \quad (\Delta\eta \gg 1) \end{aligned}$$

- Entropy becomes extensive in $\Delta\eta$!

Universal entanglement entropy density

- for very early times “Hubble” expansion rate dominates over masses and interactions

$$H = \frac{1}{\tau} \gg M = \frac{q}{\sqrt{\pi}}, m$$

- theory dominated by free, massless fermions
- universal entanglement entropy density

$$\frac{dS}{d\Delta\eta} = \frac{c}{6}$$

with conformal charge c


- for QCD in 1+1 D (gluons not dynamical, no transverse excitations)

$$c = N_c \times N_f$$

- from fluctuating transverse coordinates (Nambu-Goto action)

$$c = N_c \times N_f + 2 \approx 9 + 2 = 11$$

Modular or entanglement Hamiltonian 1



- conformal field theory
- hypersurface Σ with boundary on the intersection of two light cones
- reduced density matrix [Casini, Huerta, Myers (2011), Arias, Blanco, Casini, Huerta (2017), see also Candelas, Dowker (1979)]

$$\rho_A = \frac{1}{Z_A} e^{-K}, \quad Z_A = \text{Tr} e^{-K}$$

- modular or entanglement Hamiltonian K

Modular or entanglement Hamiltonian 2

- modular or entanglement Hamiltonian is **local expression**

$$K = \int_{\Sigma} d\Sigma_{\mu} \xi_{\nu}(x) T^{\mu\nu}(x).$$

- energy-momentum tensor $T^{\mu\nu}(x)$ of excitations
- vector field


$$\xi^{\mu}(x) = \frac{2\pi}{(q-p)^2} [(q-x)^{\mu}(x-p)(q-p) \\ + (x-p)^{\mu}(q-x)(q-p) - (q-p)^{\mu}(x-p)(q-x)]$$

end point of future light cone q , starting point of past light cone p

- inverse temperature and fluid velocity

$$\xi^{\mu}(x) = \beta^{\mu}(x) = \frac{u^{\mu}(x)}{T(x)}$$

Modular or entanglement Hamiltonian 3





- for $\Delta\eta \rightarrow \infty$: fluid velocity in τ -direction, τ -dependent temperature

$$T(\tau) = \frac{\hbar}{2\pi\tau}$$

- **Entanglement between different rapidity intervals alone leads to local thermal density matrix at very early times !**
- Hawking-Unruh temperature in Rindler wedge $T(x) = \hbar c / (2\pi x)$

Particle production in massive Schwinger model

[ongoing work with Lara Kuhn, Jürgen Berges]



- for expanding strings
- asymptotic particle number depends on $g \sim m/q$
- exponential suppression for large fermion mass $g \gg 1$

$$\frac{N}{\Delta\eta} \sim e^{-0.55 \frac{m}{q} + 7.48 \frac{q}{m} + \dots} = e^{-0.55 \frac{m}{\sqrt{2}\sigma} + 7.48 \frac{\sqrt{2}\sigma}{m} + \dots}$$

Wigner distribution and entanglement

- Classical field approximation usually based on non-negative Wigner representation of density matrix
- leads for many observables to classical statistical description
- can nevertheless show entanglement and pass Bell test for “improper” variables where Weyl transform of operator has values outside of its spectrum [Revzen, Mello, Mann, Johansen (2005)]
- Bell test violation also possible for negative Wigner distribution [Bell (1986)]

Transverse coordinates

- so far dynamics strictly confined to 1+1 dimensions
- transverse coordinates may fluctuate, can be described by Nambu-Goto action ($h_{\mu\nu} = \partial_\mu X^m \partial_\nu X_m$)

$$\begin{aligned} S_{\text{NG}} &= \int d^2x \sqrt{-\det h_{\mu\nu}} \{-\sigma + \dots\} \\ &\approx \int d^2x \sqrt{g} \left\{ -\sigma - \frac{\sigma}{2} g^{\mu\nu} \partial_\mu X^i \partial_\nu X^i + \dots \right\} \end{aligned}$$

- two additional, massless, bosonic degrees of freedom corresponding to transverse coordinates X^i with $i = 1, 2$

Temperature and entanglement entropy

- for conformal fields, entanglement entropy has also been calculated at non-zero temperature.
- for static interval of length L [Korepin (2004); Calabrese, Cardy (2004)]

$$S(T, l) = \frac{c}{3} \ln \left(\frac{1}{\pi T \epsilon} \sinh(\pi LT) \right) + \text{const}$$


- compare this to our result in expanding geometry

$$S(\tau, \Delta\eta) = \frac{c}{3} \ln \left(\frac{2\tau}{\epsilon} \sinh(\Delta\eta/2) \right) + \text{const}$$

- expressions agree for $L = \tau\Delta\eta$ (with metric $ds^2 = -d\tau^2 + \tau^2 d\eta^2$) and time-dependent temperature

$$T = \frac{1}{2\pi\tau}$$

Rapidity distribution



[open (filled) symbols: e^+e^- (pp), Grosse-Oetringhaus & Reygers (2010)]

- rapidity distribution $dN/d\eta$ has plateau around midrapidity
- only logarithmic dependence on collision energy

Experimental access to entanglement ?

- could longitudinal entanglement be tested experimentally?
- unfortunately entropy density $dS/d\eta$ not straight-forward to access
- measured in e^+e^- is the number of charged particles per unit rapidity $dN_{\text{ch}}/d\eta$ (rapidity defined with respect to the thrust axis)
- typical values for collision energies $\sqrt{s} = 14 - 206$ GeV in the range

$$dN_{\text{ch}}/d\eta \approx 2 - 4$$

- entropy per particle S/N can be estimated for a hadron resonance gas in thermal equilibrium $S/N_{\text{ch}} = 7.2$ would give

$$dS/d\eta \approx 14 - 28$$

- this is an upper bound: correlations beyond one-particle functions would lead to reduced entropy

Entanglement and QCD physics

- how strongly entangled is the nuclear wave function?
- what is the entropy of quasi-free partons and can it be understood as a result of entanglement? [Kharzeev, Levin (2017)]
- does saturation at small Bjorken- x have an entropic meaning?
- entanglement entropy and entropy production in the color glass condensate [Kovner, Lublinsky (2015); Kovner, Lublinsky, Serino (2018)]
- could entanglement entropy help for a non-perturbative extension of the parton model?
- entropy of perturbative and non-perturbative Pomeron descriptions [Shuryak, Zahed (2017)]