String Amplitudes

The interaction of n incoming/outgoing states can be written as

$$A = \sum_{\text{topologies}}$$



$$= \sum_{\text{topologies}} \int \frac{[dx][dg]}{V_{diff \times Weyl}} e^{-S_x} V_1 \cdots V_n$$

where the denominator $V_{d\times w}$ keeps into account the gauge invariance that must be modded out, and V_i is the vertex operator of the i-th in/out coming state.

We can avoid the denominator and properly quantize the theory by using the Faddeev Popov approach

Given a "preferred" metric $\hat{g}$ (flat metric) we know that there exists always a transformation ξ in Diff x Weyl such that, for each possible metric g, $g = \hat{g}^\xi = \xi(\hat{g})$ <u>locally</u>

This allows us to write

$$1 = \Delta_{FP}(g) \int d\xi \, \delta(g - \hat{g}^\xi)$$

and to replace it into the amplitude

$$A|_{\substack{\text{given} \\ \text{topology}}} = \int \frac{[dx\, dg\, d\xi]}{V_{d\times w}} \delta(g - \hat{g}^\xi) e^{-S_x} V_1 - V_n \Delta_{FP}(g)$$

We can then integrate in $d\varphi$

$$A_{g.f.} = \int \frac{[dx\, d\xi]}{Diff \times Weyl} e^{-S_x[\hat{\varphi}^\xi]} V_1 - V_n \Delta_{FP}(\hat{\varphi}^\xi)$$

moreover, since S_x is ξ-invariant, and so is Δ_{FP} (check)

$$A_{g.f.} = \int [dx]\, e^{-S_x[\hat{\varphi}]} V_1 - V_n \Delta_{FP}(\hat{\varphi}) \times \underbrace{\int \frac{d\xi}{Diff \times Weyl}}_{=1}$$

<u>Subtleties</u> In order that the "change" of variable $d\varphi = d\xi \left(\frac{\partial \varphi}{\partial \xi}\right)$, of which $\frac{\partial \varphi}{\partial \xi} \sim \Delta_{FP}$, be well defined we need a one-to-one map, i.e.

$$\forall \varphi \; \exists ! \, \xi \mid \hat{\varphi}^\xi = \varphi \quad \varphi \underline{\text{Global}}$$

1) <u>Local → Global</u>

$\Rightarrow \exists$ in general $\varphi, \varphi' \mid \nexists \xi \mid \varphi^\xi = \varphi'$

2) $\underline{\underline{\exists !}}$ $\Rightarrow \exists$ in general $\xi \mid \varphi^\xi = \varphi$

ξ is called conformal Killing vector and is part of the "gauge group" ckg

Torus case

(1) It is always possible to bring g to $\eta_{\mu\nu} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ via Diff × Weyl but the boundary conditions are then non trivial





more precisely, given

$$z = \sigma_1 + i\sigma_2$$

$$z \sim z + (m + nz)$$

and the statement is that it is not possible to reabsorb z via Diff × Weyl without a modification

$$\eta_{ab} \to g_{ab}(z)$$

Choice Toller trivial b.c.

$\Rightarrow (\sigma_1, \sigma_2) \sim (\sigma_1 + m + z_1 n, \sigma_2 + z_2 n)$

let $\sigma_1' = \sigma_1 - \sigma_2 \frac{z_1}{z_2}$

$\sigma_2' = \frac{\sigma_2}{z_2}$

$(\sigma_1', \sigma_2') \sim (\sigma_1' + m + z_1 n - z_1 n, \sigma_2' + n)$

$\Rightarrow$ cvd

but $ds^2 = d\sigma_1^2 + d\sigma_2^2 = |d\sigma_1' + d\sigma_2' z|^2$

$= g_{ab} d\sigma'^a d\sigma'^b \qquad g = \begin{pmatrix} 1 & z_1 \\ z_1 & |z|^2 \end{pmatrix}$

Consequences

A single fiducial metric $\hat{g}$ is no more sufficient since $\mathcal{O}_{\hat{g}} = \{ \hat{g}^\xi, \xi \in \text{diff-weil} \}$

$\mathcal{O}_{\hat{g}} \neq$ set of all metric

Statement $\hat{g}(z)$ is indeed sufficient

$\int dz \; \mathcal{O}_{\hat{g}(z)} =$ set of all metrics

Then Replace $1 = \Delta_{FP}(g) \int d\xi \, \delta(g - \hat{g}^\xi)$

with

$1 = \Delta_{FP}(g) \int dz \, \delta(g - \hat{g}^{\xi(z)})$

Generalization Torus $\to$ a single complex Teichmüller Modulus

Other manifolds $\to$ more real moduli

$1 = \Delta_{FP}(g) \int \prod_{i=1}^{n_e} dt^i \, \delta(g - \hat{g}^{\xi(t_i)})$

(2) $\xi: \sigma^a \longmapsto \sigma^a + \alpha^a$

is a perfectly well-defined Diff transformation

but $\xi: \eta_{\mu\nu} \longmapsto \eta_{\mu\nu}$

Statement: on the torus only these two transformations, parametrized by α^1 and α^2, defines the continuous part of the CKG

Solution: insert a σ-dependence in Δ_{FP}

$$1 = \Delta_{FP}[\sigma, g] \int \prod_i dt^i \int d\xi\, \delta(g - \hat{g}^\xi(t^i))\, \delta(\sigma_1 - \hat{\sigma}_1^\xi) \times \delta(\sigma_2 - \hat{\sigma}_2^\xi)$$

the problem is solved since, given

$\hat{\xi} \mid g = \hat{g}\hat{\xi}$, $\exists!\ \xi_{CKG} \mid \sigma = \hat{\sigma}^{\hat{\xi}\cdot\xi_{CKG}}$

Generalization

It is possible to reabsorb the ambiguity of a CKG of real dimension n_K by introducing n_K Dirac deltas

$\delta(\sigma_i - \hat{\sigma}_i^\xi)$

in the definition of Δ_{FP}

$$1 = \Delta_{FP}[\sigma_i, g] \int_F \prod_{i=1}^{n_K} dt^i \int d\xi\, \delta(g - \hat{g}^\xi) \prod_{i=1}^{n_K} \delta(\sigma_i - \hat{\sigma}_i^\xi)$$

Integration region for the moduli

$\Rightarrow$ Torus case

In principle $F \subset \mathbb{R}^2$, let's see the details

take the parametrization

$$(\sigma_1, \sigma_2) \sim (\sigma_1 + m + \tau_1 n, \sigma_2 + \tau_2 n)$$

$$ds^2 = d\sigma_1^2 + d\sigma_2^2$$

def $\omega = \sigma_1 + i\sigma_2$

$\Rightarrow ds^2 = d\omega\, d\bar{\omega}$

$$\omega \sim \omega + m + \tau n$$

Now (1) replace τ with $\tau + 1$ (T)

$\Rightarrow$ same b.c. in $m' = m + n$, $n' = n$

(2) replace τ with $-1/\tau$ (S)

$$\omega \sim \omega + m - \frac{n}{\tau}$$

redefine $\omega' = \tau\omega$

$\rightarrow \omega' \sim \omega' + m\tau - n \Rightarrow$ same as before

($ds^2 = d\omega' d\bar{\omega}'$ are Weyl inv.)

$\Rightarrow F = \mathbb{R}^2 /$ Group Generated by T and S

Statement (1) T, S generate (span) $SL(2,\mathbb{Z})$ of transformations

$$\tau' = \frac{a\tau + b}{c\tau + d} \qquad a, b, c, d \in \mathbb{Z}$$

$$ad - bc = 1$$

(2) $F \Rightarrow \operatorname{Re}\tau \in [-\tfrac{1}{2}, \tfrac{1}{2}]$

$|\tau| \geq 1$ (?, >?)

$$1 = \Delta_{FP}(g,G)\int\prod_{i=1}^{ne} dt^i \int d\xi\ \delta(g-\hat{g}(t)^\xi)\prod_{a=1}^{nu}\delta(G^a-\hat{G}^{a\xi})$$

$$\Delta_{FP}^{-1}(g,G) = \int\prod_{i=1}^{ne} dt \int d\xi\ \delta(g-\hat{g}(t)^\xi)\prod_{a=1}^{nu}\delta(G^a-\hat{G}^{a\xi})$$

now $\exists!$ zero but (last subtlety)

notice that we fixed all continuous symmetries; $\exists$ a last discrete ch. symm.

TORUS $\quad G^a \to -G^a$

To fix it as it is enough to remember that there are $n_R = 2$ zeros, and to expand around just one of them

$$\Delta_{FP}^{-1}(g,G) = n_R\int\prod_{i=1}^{ne} dt^i d\xi\ \delta(g-\hat{g}(t)^\xi)\prod_{a=1}^{nu}\delta(G^a-\hat{G}^{a\xi})$$

$$d\xi = d\omega\, d\epsilon$$

$$g-\hat{g}(t)^\xi = \delta g\Big|_{\sim \text{zero}} = (2\omega - D_c\epsilon^c)\hat{g}_{ab} - (D_a\epsilon_b + D_b\epsilon_a - 2\epsilon^c)\hat{g}_{ab} + t^i\partial_{t^i}\hat{g}_{ab}$$

$$\Delta_{FP}^{-1}(g,G) = n_R\int\prod_{i=1}^{ne} dt\, d\omega\, d\epsilon\ \delta\left((2\omega - D_c\epsilon^c)\hat{g}_{ab} - 2P\epsilon_{ab} + t^i\partial_{t^i}\hat{g}_{ab}\right)\prod_{a=1}^{2}\delta(\epsilon^a)$$

$\Leftarrow$ Specific for the Torus

exponentiate the δ's

$$\Delta_{FP}^{-1}(\hat{g},\hat{G}) = n_R\int\prod_{i=1}^{ne} dt^i d\omega\, d\epsilon^a\, d\beta^{ab}\, dx_a$$

$$\exp\left\{2\pi i\int d^2\sigma\sqrt{g}\,\beta^{ab}\left[(2\omega - D_c\epsilon^c)\mathbb{1}_{ab} + 2P\epsilon_{ab} + t^i\partial_{t^i}\hat{g}_{ab}\right]\right\}$$

$$\exp\left\{2\pi i\,\epsilon^a x_a\right\}$$

Integrate $dw \rightarrow \Pi^{ab}$ traceless $(\Pi^a{}_a = 0)$

$$\Delta_{FP}^{-1}(\hat{g},\hat{c}) = n_R \prod_{i=1}^{ul} \int dt^i \, d\epsilon^a \, d\Pi^{ab} \, dx_a$$

$$\exp\left\{ 2\pi i \int d^2\sigma \sqrt{\hat{g}} \; \Pi^{ab}\left[2P\hat{e}_{ab} - t^i \partial_t i_g \right] \right\}$$

$$\exp\left\{ 2\pi i \epsilon \cdot x \right)$$

$\rightarrow$ It is just a determinant of a new modified operator

$\Rightarrow$ Pass to Grassmann variables to "invert it"

$$\epsilon^a \rightarrow c^a$$
$$\Pi_{ab} \rightarrow b_{ab}$$
$$x \rightarrow \eta$$
$$t \rightarrow \xi$$

$$\Delta_{FP}(\hat{g},\hat{c}) = \frac{1}{n_R} \int [db\,dc] \, d\xi^i d\eta_a \; e^{-\int d^2\sigma \sqrt{g}\left[b^{ab}(P c)_{ab} + \xi^i \partial_t \eta \right)]} \; e^{\eta_a c^a(\hat{\sigma})}$$

integrate in ξ and η

$$\Delta_{FP}(\hat{g},\hat{c}) = \frac{1}{n_R} \int [db\,dc] \, e^{-S^{gh}} \times \prod_{i=1}^{ue} \left[\int d^2\sigma \sqrt{g} \; b^{ab} \partial_t i_{ab} \right] \times \prod_{a=1}^{2} c^a(\hat{\sigma})$$

Torus amplitude

$$A_{torus} = \frac{\int [dx][d\varphi]}{Vol_{Diff\times Weyl}} \int_{\tilde{F}} d\tau_1 d\tau_2 \; \Delta_{FP}(G,\varphi)$$

$$\times \int d\xi \; \delta(\varphi - \hat{\varphi}(\xi)) \, \delta(G_1 - \hat{G}_1^{\xi}) \, \delta(G_2 - \hat{G}_2^{\xi}) \, e^{-S_X[\varphi]} \prod_{i=1}^{n} V_i$$

This amplitude is <u>meaningless</u> unless an explicit G_1, G_2 integration is explicitated.

In the torus case, since the volume of the CKG is <u>finite</u> and equal to the volume of the torus itself, it is possible to introduce an integration of the form

$$1 = \frac{\int d^2G}{V_{TORUS}}$$

In other words, one fixes the CKG problem by just dividing away $V_{CKG} = V_{TORUS}$?

<u>No</u> → B-insertions in Δ_{FP} !

In the general case, knowing $V_i = \int d^2G \sqrt{g} \, V_i(G)$, we can identify the G-dependence in Δ_{FP} with the G-dep. in $n_K/2$ vertex operators.

The CKG symmetry is fixed by fixing points on the manifold + including insertions in the path integral.

In the torus case, replacing the Δ_{FP}

$$A_{torus} = \frac{1}{n_R} \int \frac{dz_1 dz_2}{V_{TORUS}} \int [dx]\,[db\,dc] \overbrace{\int d^2\sigma \sqrt{\hat{g}}\,(b^{ab})_{z_1} g_{ab}}^{B_1} \times$$

$$\times \overbrace{\int d^2\sigma \sqrt{\hat{g}}\,(b^{ab})_{z_2} g_{ab}}^{B_2}\; c^1(\hat{\sigma})\, c^2(\hat{\sigma})\; e^{-[S_X + S_{ghosts}]}$$

$n_R = 2$

$= z_2$

$V_1 - V_M$

Partition function

$$Z_{torus} = \frac{1}{2} \int_F \frac{dz_1 dz_2}{z_2} \langle B_1 B_2 c^1 c^2 \rangle_{ghosts} \times \langle 1 \rangle_{x\text{-fields}}$$

$\Rightarrow$ Relation with Coleman-Weinberg formula

$\Rightarrow$ Rôle of the ghost partition function

$\Rightarrow$ Self-regulation of UV divergences