Quantum Gravity and the Renormalization Group

Assignment 8 – Dec 13+16

Exercise 16: Fixed functions – an infinite number of couplings

Motivation: So far we have studied finite sets of beta functions. However, the non-perturbative RG flow generates infinitely many terms that have to be accounted for. The purpose of this exercise is to learn how to deal with some of the complications that are added in such a case.

For this exercise, we will briefly leave the realm of gravity, and instead study a vector field Φ^a that lives in *three* Euclidean dimensions and has an O(N) symmetry (you can think of this as a special case of a theory with N scalar fields – the "vector" he refers to the O(N) group, not to spacetime; the case N = 1 is related to the well-known Ising model). We will look at the following approximation for the effective average action:

$$\Gamma_k \simeq \int d^3x \left[\frac{1}{2} (\partial_\mu \Phi_a) (\partial^\mu \Phi^a) + V_k(\rho) \right].$$
(16.22)

For convenience, we introduced $\rho = \Phi_a \Phi^a/2$. The arbitrary k-dependent potential V_k contains in general infinitely many interaction terms. If you want, you can think of it in terms of a Taylor expansion:

$$V_k(\rho) = \sum_{n \ge 1} c_{n,k} \, \rho^n \,. \tag{16.23}$$

In this, the $c_{n,k}$ are the infinitely many k-dependent couplings.

- a) What are the mass dimensions of the field Φ^a , the potential V_k , and the coupling constants $c_{n,k}$?
- b) Given an expression for the RG flow of $V_k(\rho)$, i.e. $k\partial_k V_k(\rho)$, how would the flow of the corresponding dimensionless potential look like? *Hints:* To arrive at this, you could assume the above Taylor expansion, convert the couplings $c_{n,k}$ to their dimensionless counterparts, and then try to go back to the potential. It might also be useful to introduce a dimensionless version of ρ . If we call v_k the dimensionless version of V_k , and $\bar{\rho}$ the dimensionless version of ρ , you should get a relation of the form

$$k\partial_k V_k(\rho) = a_1 \, k\partial_k v_k(\bar{\rho}) + a_2 \, v_k(\bar{\rho}) + a_3 \, \bar{\rho} \, v'_k(\bar{\rho}) \,, \tag{16.24}$$

where $a_{1,2,3}$ are field-independent coefficients that you need to determine.

We will now study the large-N limit of this theory, in which an exact solution can be obtained. This allows us to focus on the concepts while keeping analytic control. It will be useful to focus on the derivative of the dimensionless potential, $u(\bar{\rho}) \equiv v'_k(\bar{\rho})$, where we now suppress the index k to simplify notation. This is commonly done in the literature.

Suppose that in such a limit, the k-dependence of the dimensionless derivative of the potential is given by

$$k\partial_k u(\bar{\rho}) = -2u(\bar{\rho}) + \bar{\rho} \, u'(\bar{\rho}) - \frac{1}{2} \frac{u'(\bar{\rho})}{(1+u(\bar{\rho}))^2} \,. \tag{16.25}$$

We will first focus on obtaining the fixed point solution, so we will try to solve the above differential equation for $k\partial_k u = 0$, i.e. the fixed point equation.

- c) Let us first do some structural analysis: given that the above fixed point equation for u is a first-order differential equation, how many integration constants do you expect? Does it make sense that a fixed point solution has integration constants? If not, what could fix these?
- d) Let us now try to get some more feeling for the solution. We first try a Taylor expansion. Assume that

$$u_*(\bar{\rho}) = \sum_{n=0}^N d_{n*} \,\bar{\rho}^n \,. \tag{16.26}$$

Plug this ansatz into the fixed point equation, and compute the fixed point couplings d_{n*} for a reasonable number N by expanding the equation in powers of $\bar{\rho}$ as well (do this however far you want to, maybe something like N = 4). You should be able to solve for $N - n_0$ couplings, where n_0 is the number of free integration constants found in c).

[hard question] Think about what could be done to fix the integration constants. *Hint:* A commonly used technique is to set the highest retained coupling to zero, thus imposing another condition on the fixed point couplings. One then increases N systematically and checks for the convergence of any candidate solution.

e) [hard question] The fixed point equation admits an implicit solution: instead of $u_*(\bar{\rho})$, one can solve for $\bar{\rho}(u_*)$. Show that indeed,

$$\bar{\rho} = q\sqrt{u_*} + \frac{3}{4} + \frac{3}{4}\sqrt{u_*}\arctan\sqrt{u_*} - \frac{1}{4}\frac{1}{1+u_*}$$
(16.27)

is the general solution, where q is an integration constant. *Hint:* Assume from the start that $u_*(\bar{\rho})$ can be inverted so that you can write $\bar{\rho}(u_*)$. Then express $u'_*(\bar{\rho})$ in terms of $\bar{\rho}'(u_*)$ and solve the fixed point equation for $\bar{\rho}(u_*)$.

- f) Let us now try to fix the integration constant q. For this, we can argue in the following way: the function $u_*(\bar{\rho})$ should be real and well-defined for all $\bar{\rho} \ge 0$. Does this already fix the integration constant? Also plot the resulting function $u_*(\bar{\rho})$ for this value of q.
- g) Check that your polynomial solution from d) is consistent with the solution in e). Also compute the right value of your integration constant in the Taylor solution from f).

Now that we have a fixed point function, we can compute the critical exponents! How can we do so? Let us take a step back and remember how we compute critical exponents for a finite set of beta functions. We expand the beta functions to linear order about the fixed point:

$$\vec{\beta}(\vec{d}) \simeq \underbrace{\vec{\beta}(\vec{d}_*)}_{=0} + \frac{\partial \vec{\beta}}{\partial \vec{d}} \Big|_{\vec{d}=\vec{d}_*} \left(\vec{d} - \vec{d}_*\right), \qquad (16.28)$$

where we combined our above couplings d in a vector, $\vec{d} = \{d_0, d_1, ...\}$. The solution to this linearised differential equation is

$$\vec{l} \simeq \vec{d}_* + \sum_i c_i \, \vec{e}_i \, k^{-\theta_i} \,,$$
 (16.29)

where c_i are integration constants, and the $\vec{e_i}$ are the eigenvectors of the stability matrix with eigenvalue θ_i .

Let us try to generalise this to the case of a full function. For this, we insert the linearised solution into u:

$$u(\bar{\rho}) = \vec{d} \cdot \{1, \bar{\rho}, \bar{\rho}^2, \dots\} = \underbrace{\vec{d}_* \cdot \{1, \bar{\rho}, \bar{\rho}^2, \dots\}}_{u_*(\bar{\rho})} + \sum_i c_i \left[\vec{e}_i \cdot \{1, \bar{\rho}, \bar{\rho}^2, \dots\}\right] k^{-\theta_i}.$$
 (16.30)

Here, we simply wrote the sum as a vector product of the coupling vector with the vector of all powers of $\bar{\rho}$. What is this expression? The first terms is clearly just the fixed function u_* . The second is a sum of different *eigenfunctions* that are in one-to-one correspondence with the eigenvectors of the stability matrix. This motivates the following way to compute critical exponents. Assume that

$$u(\bar{\rho}) \simeq u_*(\bar{\rho}) + \delta u(\bar{\rho}) k^{-\theta}.$$
(16.31)

We plug this into the RG equation (16.25) and expand to linear order in perturbations δu . This gives a *linear* differential equation for the perturbations δu , and θ plays the role of an eigenvalue of a differential operator. We will try to put this into practice now.

h) [hard question] Show that the eigenvalue equation for the linear perturbation δu reads

$$-\theta \delta u(\bar{\rho}) = -2\delta u(\bar{\rho}) + \bar{\rho}\,\delta u'(\bar{\rho}) + \frac{u'_*(\bar{\rho})}{(1+u_*(\bar{\rho}))^3}\delta u(\bar{\rho}) - \frac{1}{2}\frac{1}{(1+u_*(\bar{\rho}))^2}\delta u'(\bar{\rho})\,.$$
(16.32)

- i) **[hard question]** Use the fixed point equation to replace $u'_*(\bar{\rho})$ in terms of $u_*(\bar{\rho})$ and $\bar{\rho}$, and then perform a variable transform with the implicit fixed point solution $\bar{\rho}(u_*)$ to derive an equation for $\delta u(u_*)$.
- j) **[hard question]** Solve this differential equation for $\delta u(u_*)$. This should look rather ugly. To make it nicer, recall that δu is actually the $\bar{\rho}$ -derivative of the corresponding perturbation of the potential, δv . From this, derive the perturbations $\delta v(u_*)$. You should get

$$\delta v(u_*) = c \, u_*^{\frac{3-\theta}{2}} \,, \tag{16.33}$$

with c being (yet another) normalisation constant. Can you find an argument that would restrict (reasonable) perturbations which would also fix the "allowed" critical exponents?