

SPLIT SUPERSYMMETRY AT COLLIDERS

Tilman Plehn

Max Planck Institute for Physics, Munich

- Split and TeV scale supersymmetry
- Signals at the LHC
- Signals at the ILC
- What stays [W. Kilian, P. Richardson, TP, E. Schmidt: EPC C39]

TeV SCALE SUPERSYMMETRY: 1

Starting from data...

- ...which seem to indicate a light Higgs
- problem of light Higgs: scalar masses perturbatively unstable
quadratic divergences $\delta m_h^2 \propto g^2 \Lambda^2$
all-orders Higgs mass driven to cutoff $m_h \rightarrow \Lambda$
- ⇒ solution: counter term for exact cancellation ⇒ **artificial, ugly, fine tuned**
- ⇒ or new physics at TeV scale: **supersymmetry**
extra dimensions
little Higgs (pseudo-Goldstone Higgs)
Higgsless/composite Higgs
YourFavoriteNewPhysics...
- ⇒ all beautiful concepts and symmetries
- ⇒ in general problematic to realize at TeV scale [data seriously in the way]

Idea of supersymmetry: cancellation of divergences through statistics factor (-1)
[scalars vs. SM fermions; fermions vs. SM gauge bosons; fermions vs. SM scalars]

TEV SCALE SUPERSYMMETRY: 2

Bright side

- light fundamental Higgs by construction [data]
- 3 running gauge couplings meet — GUT gauge group [data]
- R parity — stable proton yields dark matter [data]
- 2 Higgs doublets — radiative symmetry breaking [beauty]
- local supersymmetry – including gravity? [beauty]
- **rich LHC phenomenology** [effective theory of everything short-lived]

Dark side

- unknown Susy breaking
→ masses, couplings, phases
 - flavor physics and Susy breaking
→ CKM and lepton flavor?
 - 2 Higgs doublet model
→ μ and Susy breaking? [Giudice, Masiero]
- ⇒ as many exclusive analyses as possible

		spin	d.o.f.	
gluon	G_μ	1	n-2	
→ gluino	\tilde{g}	1/2	2	Majorana
gauge bosons	γ, Z	1	2+3	
Higgs bosons	h^0, H^0, A^0	0	3	
→ neutralinos	$\tilde{\chi}_i^0$	1/2	4 · 2	Majorana
gauge bosons	W^\pm	1	2 · 3	
Higgs bosons	H^\pm	0	2	
→ charginos	$\tilde{\chi}_i^\pm$	1/2	2 · 4	Dirac
fermion	f_L, f_R	1/2	1+1	
→ sfermion	\tilde{f}_L, \tilde{f}_R	0	1+1	

[never believe us theorists when we say we know...]

TeV SCALE SUPERSYMMETRY: 3

Gauginos and higgsinos in the SUSY spectrum [Dimopoulos; Drees, Martin]

- gauginos–higgsinos mixing: $m_{\tilde{\chi}_2^0} \sim m_{\tilde{\chi}_1^+}$ or $m_{\tilde{\chi}_1^0} \sim m_{\tilde{\chi}_1^+}$ in **MSSM**

$$\begin{pmatrix} m_{\tilde{g}} & 0 & -m_Z s_W c_\beta & m_Z s_W s_\beta \\ 0 & m_{\tilde{W}} & m_Z c_W c_\beta & -m_Z c_W s_\beta \\ -m_Z s_W c_\beta & m_Z c_W c_\beta & 0 & -\mu \\ m_Z s_W s_\beta & -m_Z c_W s_\beta & -\mu & 0 \end{pmatrix} \begin{pmatrix} m_{\tilde{W}} & \sqrt{2} m_Z c_W s_\beta \\ \sqrt{2} m_Z c_W c_\beta & -\mu \end{pmatrix}$$

- heavy gluinos through **unification**: $m_{\tilde{B}, \tilde{W}, \tilde{g}}/m_{1/2} \sim 0.4, 0.8, 2.6$

[mass and coupling unification independent]

- lightest Susy partner $\tilde{\chi}_1^0, \tilde{\nu}$

\Rightarrow after dark matter data $\tilde{\chi}_1^0 \sim \tilde{B}, \tilde{W}$ [Ellis, Falk, Olive...]

PHYSICS OF SPLIT SUPERSYMMETRY: 1

Split supersymmetry [Dimopoulos, Arkani-Hamed; Giudice, Romanino]

- forget about fine tuning [Higgs will never be as bad as cosmological constant]
- remember all the good things Susy did for you [dark matter, unification from data]
- notice that scalars are evil [lepton and quark flavor, Higgs mass and LEP2]
- remember simple facts about unification [SU(5) multiplets decouple; Dawson, Georgi 1979]
- ⇒ **make all scalars heavy** [hope: $\tilde{m} \rightarrow m_{\text{GUT}}?$]
- ⇒ protect all gaugino and higgsino masses [$m_{\tilde{\chi}_i}, m_{\tilde{g}} \lesssim \text{TeV}$]


Fine tuning no excuse for multi-billion dollar experiments [trigger by popular vote of theorists?]

- gluinos and gauginos at the LHC
- gauginos and higgsinos at the ILC
- ⇒ **is it supersymmetry?**
- ⇒ **is it split?**

PHYSICS OF SPLIT SUPERSYMMETRY: 2

Heavy scalars and the Higgs mass [Giudice, Romanino; Arvantaki, Davis, Graham, Wacker]


- known leading corrections increased: $m_h \sim m_Z + G_F y_t^4 \log(m_{\tilde{t}}^2/m_t^2)$
- ⇒ large m_h for heavy stops [out of LEP2 reach]
- ⇒ not a precision observable anymore [large logarithms]
- ⇒ light Higgs is a SM Higgs boson with $m_h \gtrsim 140$ GeV



PHYSICS OF SPLIT SUPERSYMMETRY: 2


Heavy scalars and the Higgs mass [Giudice, Romanino; Arvantaki, Davis, Graham, Wacker]

- known leading corrections increased: $m_h \sim m_Z + G_{Fy_t^4} \log(m_t^2/m_t^2)$
- ⇒ large m_h for heavy stops [out of LEP2 reach]
- ⇒ not a precision observable anymore [large logarithms]
- ⇒ light Higgs is a SM Higgs boson with $m_h \gtrsim 140$ GeV



Heavy scalars and the gluino life time [Arkani-Hamed, Dimopoulos; Giudice, Romanino]



- decay through squark $\tau_{\tilde{g}} \sim \tilde{m}^4/m_{\tilde{g}}^5$
- loop-induced decays? [Toharia, Wells]
- lifetime constrained by age of universe
- $\tilde{m} \lesssim 10^{12} \text{ GeV} \ll m_{\text{GUT}}$ [PeV? Wells]
- ⇒ gluino hadronizes, decays much later
- ⇒ **long-lived gluino collider signature No.1**



PHYSICS OF SPLIT SUPERSYMMETRY: 3

Renormalization group running


- argued unification, so make Split Susy a GUT
- gauge couplings unify
- gaugino masses as well



PHYSICS OF SPLIT SUPERSYMMETRY: 3

Renormalization group running

- argued unification, so make Split Susy a GUT
- gauge couplings unify
- gaugino masses assumed to unify as well




Anomalous into Yukawa coupling

- gauginos–higgsinos mixing in MSSM:

$$\begin{pmatrix} m_{\tilde{B}} & 0 & -m_Z s_W c_\beta \equiv -\tilde{g}_d v & m_Z s_W s_\beta \equiv \tilde{g}_u v \\ 0 & m_{\tilde{W}} & m_Z c_W c_\beta \equiv \tilde{g}'_d v & -m_Z c_W s_\beta \equiv -\tilde{g}'_u v \\ -m_Z s_W c_\beta & m_Z c_W c_\beta & 0 & -\mu \\ m_Z s_W s_\beta & -m_Z c_W s_\beta & -\mu & 0 \end{pmatrix}$$

- Yukawas/gaugino–higgsino mixing fixed by Susy
- supersymmetric beta functions broken at $Q = \tilde{m}$
- **anomalous Yukawas collider signal No.2: $\tilde{g}/\tilde{g}_{\text{MSSM}} = 1 + \kappa$**



SPLIT SUSY AT THE LHC: 1

LHC production of gauginos and higgsinos

- cross sections not small $[M_j(m_{\text{GUT}}) = 120\text{GeV}; \sigma \text{ in fb from Prospino2}]$

$\tilde{g}\tilde{g}$	1710						
$\tilde{\chi}_1^- \tilde{\chi}_1^+$	2910	$\tilde{\chi}_1^- \tilde{\chi}_2^+$	73.7	$\tilde{\chi}_1^+ \tilde{\chi}_2^-$	73.7	$\tilde{\chi}_2^+ \tilde{\chi}_2^-$	604
$\tilde{\chi}_1^0 \tilde{\chi}_1^0$	49.4	$\tilde{\chi}_1^0 \tilde{\chi}_2^0$	49.7	$\tilde{\chi}_1^0 \tilde{\chi}_3^0$	409	$\tilde{\chi}_1^0 \tilde{\chi}_4^0$	0.06
		$\tilde{\chi}_2^0 \tilde{\chi}_2^0$	5.0	$\tilde{\chi}_2^0 \tilde{\chi}_3^0$	876	$\tilde{\chi}_2^0 \tilde{\chi}_4^0$	3.7
				$\tilde{\chi}_3^0 \tilde{\chi}_3^0$	1.4	$\tilde{\chi}_3^0 \tilde{\chi}_4^0$	69.6
						$\tilde{\chi}_4^0 \tilde{\chi}_4^0$	1.0
$\tilde{\chi}_1^- \tilde{\chi}_1^0$	584	$\tilde{\chi}_1^- \tilde{\chi}_2^0$	1780	$\tilde{\chi}_1^- \tilde{\chi}_3^0$	789	$\tilde{\chi}_1^- \tilde{\chi}_4^0$	78.8
$\tilde{\chi}_1^+ \tilde{\chi}_1^0$	914	$\tilde{\chi}_1^+ \tilde{\chi}_2^0$	2870	$\tilde{\chi}_1^+ \tilde{\chi}_3^0$	1310	$\tilde{\chi}_1^+ \tilde{\chi}_4^0$	138
$\tilde{\chi}_2^- \tilde{\chi}_1^0$	2.7	$\tilde{\chi}_2^- \tilde{\chi}_2^0$	55.9	$\tilde{\chi}_2^- \tilde{\chi}_3^0$	66.6	$\tilde{\chi}_2^- \tilde{\chi}_4^0$	430
$\tilde{\chi}_2^+ \tilde{\chi}_1^0$	4.5	$\tilde{\chi}_2^+ \tilde{\chi}_2^0$	97.7	$\tilde{\chi}_2^+ \tilde{\chi}_3^0$	119	$\tilde{\chi}_2^+ \tilde{\chi}_4^0$	798

- but best background rejection $m_{\ell\ell}$ gone with the wind [higgsino searches?]


What's new for LHC phenomenology?

- no squarks, sleptons for cascades [Giudice, Romanino; astro-particle: Pierce]
- stable (hadronizing) gluinos $[\tau \sim \tilde{m}^{-4} \sim 6.5\text{s} \text{ for } \tilde{m} = 10^9\text{GeV}, \text{LHC time scale } 25 \text{ ns}]$
- heavy hadrons $R_g, R_{q\bar{q}}, R_{qqq}$ [Farrar, Fayet 1978; Baer, Cheung, Gunion 1999; UKQCD 1999]
- gluinonium [Kühn, Ono 1984; Goldman, Haber 1985; Cheung]

SPLIT SUSY AT THE LHC: 2

Charged R hadrons

- many gluinos pair-produced [$\sigma \gtrsim 1$ pb]
 - charged R hadrons in tracker, calorimeter, muon chambers [Cambridge ex-th]
 - level-1 trigger without muon chamber? [25...75 ns delay]
 - effect of conversion to R baryons because of light pions? [Kraan]
- ⇒ fraction of charged R hadrons crucial
- ⇒ effective (not calculable) parameter P_{R_g}




SPLIT SUSY AT THE LHC: 2

Charged R hadrons

- many gluinos pair-produced [$\sigma \gtrsim 1$ pb]
 - charged R hadrons in tracker, calorimeter, muon chambers [Cambridge ex-th]
 - level-1 trigger without muon chamber? [25...75 ns delay]
 - effect of conversion to R baryons because of light pions? [Kraan]
- ⇒ fraction of charged R hadrons crucial
- ⇒ effective (not calculable) parameter P_{R_g}

Beyond BSM signal


- mass measurement through time of flight
- charge flipper [Kraan; Hewett, Rizzo,...]
- energy deposition: no heavy lepton



SPLIT SUSY AT THE LHC: 3

Neutral R hadrons


- jets plus missing energy [$\sim 10\%$ energy loss]
 - trigger dependent on cross section in calorimeter
 - improved in combination with charged R hadron [missing energy trigger]
 - mass measurement from gluinoonium
 - R hadron flavor physics?
- ⇒ **charged R hadrons preferable**



SPLIT SUSY AT THE LHC: 3

Neutral R hadrons


- jets plus missing energy [$\sim 10\%$ energy loss]
 - trigger dependent on cross section in calorimeter
 - improved in combination with charged R hadron [missing energy trigger]
 - mass measurement from gluinoonium
 - R hadron flavor physics?
- ⇒ **charged R hadrons preferable**



SPLIT SUPERSYMMETRY AT THE ILC: 1

Signals at the ILC


- gluinos not produced because of decoupled squarks
 - neutralino–chargino sector analysis as usual [robust with changed decay channels]
 - measurement of anomalous Yukawas [$\tilde{g}_u, \tilde{g}_d, \tilde{g}'_u, \tilde{g}'_d$ different by $\sim 10\%$]
- ⇒ (1) direct measurements of $\chi\chi H$



SPLIT SUPERSYMMETRY AT THE ILC: 1


Signals at the ILC

- gluinos not produced because of decoupled squarks
 - neutralino–chargino sector analysis as usual [robust with changed decay channels]
 - measurement of anomalous Yukawas [$\tilde{g}_u, \tilde{g}_d, \tilde{g}'_u, \tilde{g}'_d$]
- ⇒ (1) direct measurements of $\chi\chi H$ [Whizard, Smadgraph; unpromising!]
(2) indirect determination of mass matrices




Extracting parameters from neutralino/chargino sector

- 10^4 smeared measurements of six masses (and cross sections)
- 10^4 fits of M_1, M_2, μ and one or more κ_i
- LHC data alone not promising [masses only, 5% error]



Neutralinos/charginos at the ILC


- mass measurements to 0.5%
 - error propagation through 10^4 smeared pseudo-measurements
- ⇒ one κ at the time to $\lesssim 5\%$



SPLIT SUPERSYMMETRY AT THE ILC: 2

Neutralinos/charginos at the ILC

- mass measurements to 0.5%, cross sections statistical error
 - error propagation through 10^4 smeared pseudo-measurements
- ⇒ one κ at the time to $\lesssim 5\%$
- ⇒ four κ simultaneously to $\lesssim 10\%$



SPLIT SUPERSYMMETRY AT THE ILC: 2

Neutralinos/charginos at the ILC

- mass measurements to 0.5%, cross sections statistical error
- error propagation through 10^4 smeared pseudo-measurements
- ⇒ one κ at the time to $\lesssim 5\%$
- ⇒ four κ simultaneously to $\lesssim 10\%$

So can we tell it is Split Susy?

- mass measurement errors conservative
- only mass and cross section measurements yet [Sfitter-Fittino next step]

	Fit $\tan\beta$	m_i	σ_{ij}	$\Delta\kappa_U$	$\Delta\kappa_D$	$\Delta\kappa'_U$	$\Delta\kappa'_D$
ILC		•	•	0.9×10^{-2}	3×10^{-2}	1.3×10^{-2}	4×10^{-2}
ILC	•	•	•	1.2×10^{-2}	5×10^{-2}	2×10^{-2}	5×10^{-2}
ILC		•		1.1×10^{-2}	5×10^{-2}	3×10^{-2}	8×10^{-2}
ILC	•	•		1.2×10^{-2}	11×10^{-2}	4×10^{-2}	8×10^{-2}
LHC		•		2.2×10^{-1}	6×10^{-1}	2.7×10^{-1}	8×10^{-1}
ILC		•	•	1.4×10^{-2}	5×10^{-2}	3×10^{-2}	10×10^{-2}
ILC*	•	•	•	1.7×10^{-2}	9×10^{-2}	4×10^{-2}	13×10^{-2}
ILC	fix $\tan\beta = 3$	•	•	1.6×10^{-2}	4×10^{-2}	4×10^{-2}	9×10^{-2}
ILC*	$\kappa_i \neq 0$	•	•	1.4×10^{-2}	5×10^{-2}	4×10^{-2}	11×10^{-2}
ILC*	fix $\tan\beta = 5$	•	•	1.6×10^{-2}	7×10^{-2}	4×10^{-2}	14×10^{-2}

⇒ **anomalous Yukawas promising at ILC**

Showcase for state-of-the-art LHC phenomenology: Split Supersymmetry

- interesting phenomenology
- LHC: R hadrons observable with mass measurement
- ILC: anomalous weak-ino Yukawas accessible

What stays

- exotic heavy hadrons visible at LHC [trigger issues]
- why did we always assume MSSM-type ino Yukawas? [missed Susy test]