

SUPERSYMMETRY AT THE LHC

Tilman Plehn

Max–Planck–Institut für Physik, München

- TeV scale supersymmetry
- Signals at LHC: Prospino2, Smadgraph
- Measurements at LHC (+ILC): Sfitter
- LHC phenomenology at work: split supersymmetry

TEV SCALE SUPERSYMMETRY: 1

Starting from data...

- ...which seem to indicate a light Higgs
- problem of light Higgs: scalar masses perturbatively unstable
quadratic divergences $\delta m_h^2 \propto g^2 \Lambda^2$
all-orders Higgs mass driven to cutoff $m_h \rightarrow \Lambda$
- ⇒ solution: counter term for exact cancellation ⇒ **artificial, unmotivated, ugly**
- ⇒ or new physics at TeV scale: **supersymmetry**
extra dimensions
little Higgs (pseudo–Goldstone Higgs)
topcolor (composite Higgs)
YourFavoriteNewPhysics...
- ⇒ all beautiful concepts und symmetries
- ⇒ in general problematic to realize at TeV scale [data seriously in the way]

Idea of supersymmetry: cancellation of divergences through statistics factor (-1)
[scalars vs. SM fermions; fermions vs. SM gauge bosons; fermions vs. SM scalars]

Bright side

- 3 running gauge couplings meet — GUT gauge group
- 2 Higgs doublets — radiative symmetry breaking
- R parity — stable proton yields dark matter
- local supersymmetry – including gravity?
- rich LHC phenomenology — no nasty surprises [effective theory of everything]

Dark side

- unknown SUSY breaking
 - masses, scalar couplings, phases...
 - e.g. hierarchical spectrum? [Split SUSY]
 - flavor physics and SUSY breaking
 - CKM and lepton flavor?
 - 2 Higgs doublet model
 - μ parameter and SUSY breaking?
- ⇒ as many as exclusive analyses as possible

		spin	d.o.f.	
quark	\tilde{q}_L, \tilde{q}_R	1/2	1+1	6 flavors
→ squark	\tilde{q}_L, \tilde{q}_R	0	1+1	
gluon	$\tilde{G}\mu$	1	n-2	Majorana
→ gluino	g	1/2	2	
gauge bosons	$\tilde{\gamma}, \tilde{Z}$	1	2+3	
Higgs bosons	$\tilde{h}^0, \tilde{H}^0, \tilde{A}^0$	0	3	
→ neutralinos	$\tilde{\chi}_i^0$	1/2	4 · 2	Majorana
gauge bosons	\tilde{W}^\pm	1	2 · 3	
Higgs bosons	\tilde{H}^\pm	0	2	
→ charginos	$\tilde{\chi}_i^\pm$	1/2	2 · 4	Dirac

Structures in the SUSY spectrum [Drees, Martin]

- gauginos–higgsinos mixing: $m_{\tilde{\chi}_2^0} \sim m_{\tilde{\chi}_1^+}$ or $m_{\tilde{\chi}_1^0} \sim m_{\tilde{\chi}_1^+}$ in **MSSM**

$$\begin{pmatrix} m_{\tilde{B}} & 0 & -m_Z s_w c_\beta & m_Z s_w s_\beta \\ 0 & m_{\tilde{W}} & m_Z c_w c_\beta & -m_Z c_w s_\beta \\ -m_Z s_w c_\beta & m_Z c_w c_\beta & 0 & -\mu \\ m_Z s_w s_\beta & -m_Z c_w s_\beta & -\mu & 0 \end{pmatrix} \begin{pmatrix} m_{\tilde{W}} & \sqrt{2} m_w s_\beta \\ \sqrt{2} m_w c_\beta & -\mu \end{pmatrix}$$

- stop and sbottom mixing in **MSSM**

$$\begin{pmatrix} m_Q^2 + m_t^2 + \left(\frac{1}{2} - \frac{2}{3}s_w^2\right) m_Z^2 c_{2\beta} & -m_t (A_t + \mu \cot \beta) \\ -m_t (A_t + \mu \cot \beta) & m_U^2 + m_t^2 + \frac{2}{3}s_w^2 m_Z^2 c_{2\beta} \end{pmatrix}$$

- heavy gluinos and squarks through **unification**: $m_{\tilde{B}, \tilde{W}, \tilde{g}} / m_{1/2} \sim 0.4, 0.8, 2.6$
 $m_{\tilde{\ell}, \tilde{q}} / m_{1/2} \sim 0.7, 2.5$ [$m_0 \ll m_{1/2}$]

[mass and coupling unification independent]

Supersymmetric parameter conventions

- comparison of specialized codes crucial [remember: e.g. Comphep–Pythia–Isajet]
- ⇒ fix SUSY conventions once for all
 - soft breaking parameters [e.g. $\pm A_t$]
 - scale dependence of couplings, masses [e.g. $m(q = \text{TeV}, v, m_t)$?]
 - definitions of mass matrixes, mixing angles [e.g. $\tilde{t}_{L,R}$ up or down?]

SUSY Les Houches Accord [P. Skands et al.]

- spectrum generators: SoftSusy, SPheno, FeynHiggs,...
- multi-purpose Monte Carlos: Pythia, Herwig, Sherpa
- matrix element generators: Whizard, Smadgraph
- NLO cross sections: Prospino2
- NLO decay rates: Sdecay
- SUSY parameter extraction: Fittino, Sfitter
- dark matter: Micromegas
- ⇒ **fixed parameter convention and read-write format** [list to be extended]


SUSY SIGNALS AT LHC: 1

Supersymmetry at the LHC

- (1) **possible discovery** — signals for new physics, exclusion of parameter space
 - (2) **measurements** — masses, cross sections, decays
 - (3) **parameter studies** — MSSM Lagrangean, SUSY breaking
- ⇒ at least 10% precision to be matched at LHC [theorist's nightmare, yet unsolved]

Hadron collider observables with errors

- ★ masses from σ_{tot}
 - ★ branching fractions from σ_{tot}
 - renormalization scale from $\alpha_s, y_{b,t}$
 - factorization scale from pdf's
 - perturbative series $N_c \alpha_s / \pi \sim 10\%$
 - finite terms [LO-NLO-NNLO: DY, Higgs]
- ⇒ NLO errors: 15...40 % for SUSY particles



SUSY SIGNALS AT LHC: 2


Prospino2: NLO cross sections for Tevatron and LHC

- all two-particle SUSY production channels included
- download from Prospino2 page: <http://pheno.physics.wisc.edu/~plehn>
- extended version beyond Prospino2: $pp \rightarrow SS^*, tH^- \dots$

[thanks to: W. Beenakker, R. Höpker, M. Krämer, M. Spira, P. Zerwas]

SUSY signals included

- jets and E_T : $pp \rightarrow \tilde{q}\tilde{q}^*, \tilde{g}\tilde{g}, \tilde{q}\tilde{g}$
- bottoms and E_T : $pp \rightarrow \tilde{b}_1\tilde{b}_1^*$
- like sign dileptons: $pp \rightarrow \tilde{g}\tilde{g}$
 $[\tilde{g} \rightarrow \tilde{u}\bar{u} \rightarrow \tilde{\chi}_1^+ d\bar{u} \text{ or c.c.}]$
- tri-leptons: $pp \rightarrow \tilde{\chi}_2^0 \tilde{\chi}_1^-$
 $[\tilde{\chi}_2^0 \rightarrow \tilde{\ell}\bar{\ell} \rightarrow \tilde{\chi}_1^0 \ell\bar{\ell}; \tilde{\chi}_1^- \rightarrow \tilde{\chi}_1^0 \ell\bar{\nu}]$
- funny tops: $pp \rightarrow \tilde{t}_1\tilde{t}_1^*$




SUSY MEASUREMENTS AT LHC: 1

SUSY spectra from cascade decays

- decay $\tilde{g} \rightarrow \tilde{q}\bar{q} \rightarrow \tilde{\chi}_2^0 q\bar{q} \rightarrow \mu^+ \mu^- q\bar{q} \tilde{\chi}_1^0$ [better not via Z or to τ]
- cross sections some 100 pb [more than 3×10^5 events]
- thresholds & edges [Hinchliffe, Paige...; Cambridge ex-th]
critical: enough thresholds and edges available?


$$\text{classical } m_{\ell\ell}^2 < (m_{\tilde{\chi}_2^0}^2 - m_{\tilde{\ell}}^2)(m_{\tilde{\ell}}^2 - m_{\tilde{\chi}_1^0}^2)/m_{\tilde{\ell}}^2$$

\Rightarrow detector resolution, calibration, systematic errors?



Side remark: problem in decay studies

- typical cuts: $p_{T,j} > 150, 100, 50, 50$ GeV
- (a) cuts on $p_{T,j}$ hierarchy?
(b) combinatorics through jet radiation?
- \Rightarrow matrix elements for SUSY + hard jets
- \Rightarrow **Smadgraph** [Hagiwara, Kanzaki, TP, Rainwater, Stelzer]



SUSY MEASUREMENTS AT LHC: 2

Theorist's point of view

- measured masses, cross sections, decays secondary
 - parameters in SUSY Lagrangean from measurements
- ⇒ SUSY breaking parameters at TeV (or higher) scale

Warmup exercise: Sugra top-down fit

- fit including all errors

[Allanach et al; Jack & Jones]

abs. errors	SPS1a	Δ at LHC		Δ at LC		Δ at LHC+LC	
		stat	stat+theo	stat	stat+theo	stat	stat+theo
m_0	100	4.0	4.7	0.09	0.6	0.08	0.6
$m_{1/2}$	250	1.8	2.6	0.13	0.6	0.11	0.5
$\tan\beta$	10	1.3	3.5	0.14	0.3	0.14	0.4
A_0	-100	31.8	32.4	4.43	8.5	4.23	12.6

- spectrum from Suspect [Djouadi, Kneur]
- fit Suspect and Softsusy [Allanach]

LHC	Suspect	Δ	Softsusy	Δ
m_0	100.00	4.7	97.9	4.6
$m_{1/2}$	250.00	2.7	252.5	2.9
$\tan\beta$	10.00	3.5	11.6	3.6
A_0	-99.96	32.4	14.7	58.9

LHC+LC				
m_0	100.0	0.59	98.4	0.7
$m_{1/2}$	249.99	0.49	254.3	0.8
$\tan\beta$	9.99	0.44	7.3	0.3
A_0	-100.1	12.6	902.0	18

⇒ best way to estimate theory errors?

SUSY MEASUREMENTS AT LHC: 3

SUSY parameters from observables

- parameters: weak-scale MSSM Lagrangean
- measurements: masses [Suspect, Softsusy, FeynHiggs...]
branching fractions [MSMLib, Sdecay]
cross sections [Prospino, MSMLib],...
- errors: general correlation, statistics & systematics & theory
- problem in grid: huge phase space, local minimum?
problem in fit: domain walls, starting values, global minimum?

SFitter [Lafaye, TP, D. Zerwas, also Fittino]

- (1) grid for closed subset
(2) fit of remaining parameters
(3) complete fit

⇒ LHC+LC with no assumptions

	LHC	LC	LHC+LC	SPS1a
$\tan\beta$	10.22 ± 9.1	10.26 ± 0.3	10.06 ± 0.2	10
M_1	102.45 ± 5.3	102.32 ± 0.1	102.23 ± 0.1	102.2
M_3	578.67 ± 15	fix 500	588.05 ± 11	589.4
$M_{\tilde{\tau}_L}$	fix 500	197.68 ± 1.2	199.25 ± 1.1	197.8
$M_{\tilde{\tau}_R}$	129.03 ± 6.9	135.66 ± 0.3	133.35 ± 0.6	135.5
$M_{\tilde{\mu}_L}$	198.7 ± 5.1	198.7 ± 0.5	198.7 ± 0.5	198.7
$M_{\tilde{q}_3_L}$	498.3 ± 110	497.6 ± 4.4	521.9 ± 39	501.3
$M_{\tilde{t}_R}$	fix 500	420 ± 2.1	411.73 ± 12	420.2
$M_{\tilde{b}_R}$	522.26 ± 113	fix 500	504.35 ± 61	525.6
A_τ	fix 0	-202.4 ± 89.5	352.1 ± 171	-253.5
A_t	-507.8 ± 91	-501.95 ± 2.7	-505.24 ± 3.3	-504.9
A_b	-784.7 ± 35603	fix 0	-977 ± 12467	-799.4

SPLIT SUSY AT COLLIDERS: 1

Split Supersymmetry [Dimopoulos, Arkani-Hamed; Giudice, Romanino; Wells; Drees]

- forget about fine tuning [Higgs will never be as bad as cosmological constant]
 - remember all the good things SUSY did for you [dark matter, unification]
- ⇒ make all scalars heavy [hope: $\tilde{m} \rightarrow m_{\text{GUT}}$?]
- ⇒ protect all gaugino and higgsino masses [$m_{\tilde{\chi}_i}, m_{\tilde{g}} \lesssim \text{TeV}$]

What's new for phenomenology?

- no squarks, sleptons for colliders, astro-particle physics [Giudice, Romanino; Pierce]
- no cascade decays
- stable (hadronizing) gluinos [$\tau \sim \tilde{m}^{-4} \sim 6.5\text{s}$ for $\tilde{m} = 10^9\text{GeV}$]
- heavy hadrons $R_g, R_{q\bar{q}}, R_{qqq}$ [Farrar, Fayet; Baer, Cheung, Gunion; UKQCD; Kraan]
- renormalization group running without scalars [e.g. differentino Yukawa couplings by $\lesssim 20\%$]



Collider tests

- (1) Is it supersymmetry?
- (2) Is it split?

SPLIT SUSY AT COLLIDERS: 2

Split SUSY at the LHC [Kilian, TP, Richardson, Schmidt]

- neutralinos, charginos like in MSSM, poor precision [Prospino2]
- many gluinos pair-produced [$\sigma \gtrsim 1 \text{ pb}$, Prospino2]
- gluinonium $\tilde{g}\tilde{g} \rightarrow jj$ [Kühn, Ono; Goldman, Haber; CMS; reach $\sim \text{TeV?}$]
- neutral R hadrons missing \rightarrow missing energy signal
- charged R hadrons in tracker, calorimeter, muon chambers [Cambridge ex-th]
- mass measurement through time of flight tracker–muon chamber




SPLIT SUSY AT COLLIDERS: 3

Split Supersymmetry at the ILC [Kilian, TP, Richardson, Schmidt]

- gluinos not produced because of decoupled squarks
 - neutralino–chargino sector analysis as usual [robust towards decay channels]
 - anomalous Yukawas \equiv off-diagonal mass matrix entries [$g s_\beta, g c_\beta, g' s_\beta, g' c_\beta$]
- \Rightarrow (1) direct measurements of $\chi\chi h$ [[Whizard](#), [Smadgraph](#) \rightarrow distinctly unpromising]
(2) indirect determination of mass matrices [poor man's [Sfi ttbar](#)]

Indirect determination

- errors crucial [0.5 % error on masses at ILC]
 - 10^4 smeared pseudo-measurements to extract parameters [no Gaussian assumed]
- \Rightarrow analytic inversion impossible, fit instead



SPLIT SUSY AT COLLIDERS: 4

Error on anomalous Yukawa couplings

	Fit $\tan\beta$	m_i	σ_{jj}	Δ_{κ_u}	Δ_{κ_d}	$\Delta_{\kappa'_u}$	$\Delta_{\kappa'_d}$
ILC		•	•	0.9×10^{-2}	3×10^{-2}	1.3×10^{-2}	4×10^{-2}
ILC	•	•	•	1.2×10^{-2}	5×10^{-2}	2×10^{-2}	5×10^{-2}
ILC		•		1.1×10^{-2}	5×10^{-2}	3×10^{-2}	8×10^{-2}
ILC	•	•		1.2×10^{-2}	11×10^{-2}	4×10^{-2}	8×10^{-2}
LHC		•		2.2×10^{-1}	6×10^{-1}	2.7×10^{-1}	8×10^{-1}
ILC		•	•	1.4×10^{-2}	5×10^{-2}	3×10^{-2}	10×10^{-2}
ILC*	•	•	•	1.7×10^{-2}	9×10^{-2}	4×10^{-2}	13×10^{-2}
ILC	fix $\tan\beta = 3$	•	•	1.6×10^{-2}	4×10^{-2}	4×10^{-2}	9×10^{-2}

Verdict

- LHC: stable R hadrons, charginos and neutralinos
- ILC: anomalous Yukawa couplings
- IceCube: one event per year for low-mass R hadrons [Hewett, Lillie, Mazip, Rizzo]
- Pierre Auger: few events for $\tilde{m} < 10^{11}$ GeV [Anchordoqui, Goldberg, Nunez]
- ⇒ split supersymmetry identifiable at combination of colliders
- ⇒ what stays: exotic heavy hadrons visible at LHC
why did we ever assume MSSM-typeino Yukawas?

OUTLOOK


Theory effort for SUSY at the LHC well advanced

- cascade reconstruction with great promise
 - total cross sections available to NLO [Propino2]
 - automatic matrix element generators being tested [Smadgraph, Whizard, Sherpa]
 - parameter extraction tools in use [Sfitter, Fittino]
- ⇒ We can do errors now!


Showcase: Split Supersymmetry

- interesting phenomenology
- LHC: R hadrons observable with mass measurement
- ILC: anomalous weak-ino Yukawas accessible




APPENDIX: 1



APPENDIX: 2



APPENDIX: 3



Prospino2: NLO cross sections for Tevatron and LHC

- all two-particle SUSY production channels included
- download from Prospino2 page: <http://pheno.physics.wisc.edu/~plehn>
- extended version beyond Prospino2: $pp \rightarrow SS^*, tH^- \dots$

[thanks to: W. Beenakker, R. Höpker, M. Krämer, M. Spira, P. Zerwas]

SUSY signals included

- jets and E_T : $pp \rightarrow \tilde{q}\tilde{q}^*, \tilde{g}\tilde{g}, \tilde{q}\tilde{g}$
 $[\tilde{g} \rightarrow \tilde{u}\tilde{u} \rightarrow \tilde{\chi}_1^+ d\bar{u} \text{ or c.c.}]$
- bottoms and E_T : $pp \rightarrow \tilde{b}_1\tilde{b}_1^*$
- like sign dileptons: $pp \rightarrow \tilde{g}\tilde{g}$
- tri-leptons: $pp \rightarrow \tilde{\chi}_2^0\tilde{\chi}_1^-$
 $[\tilde{\chi}_2^0 \rightarrow \ell\bar{\ell} \rightarrow \tilde{\chi}_1^0\ell\bar{\ell}; \tilde{\chi}_1^- \rightarrow \tilde{\chi}_1^0\ell\bar{\nu}]$
- funny tops: $pp \rightarrow \tilde{t}_1\tilde{t}_1^*$

