

New Methods for New Physics

Tilman Plehn

MPI für Physik & University of Edinburgh

DESY, 12/2006

Outline

Supersymmetry

LHC Signals

Masses

Spins

Parameters

Statistics

TeV-scale supersymmetry

Supersymmetric signatures at LHC

New physics mass measurements


New physics spin measurements

Supersymmetric parameter studies

Statistics and phenomenology

TeV-scale supersymmetry: 1

Starting from data...



- ...which seems to indicate a light Higgs [e-w precision data]
- problem of light Higgs: mass driven to cutoff of effective Standard Model:

$$\delta m_H^2 \propto g^2 (2m_W^2 + m_Z^2 + m_H^2 - 4m_t^2) \Lambda^2$$
- ⇒ easy solution: counter term to cancel loops ⇒ **artificial, unmotivated, ugly**
- ⇒ or new physics at TeV scale:
 - supersymmetry**
 - extra dimensions
 - little Higgs (pseudo-Goldstone Higgs)
 - Higgsless, composite Higgs, TopColor, YourFavoriteNewPhysics...
- ⇒ typically cancellation by new particles or discussing away high scale
- ⇒ beautiful concepts, but problematic at TeV scale [data seriously in the way]
- ⇒ **new physics models in baroque state**

Idea of supersymmetry:

cancellation of divergences through statistics factor (-1)

[SM fermions to scalar; SM gauge bosons to fermions; SM scalars to fermions]

TeV-scale supersymmetry: 2

SUSY breaking: (yet) unobserved partners heavy

- link to **BSM dark matter** [Falk,...]
- link to **BSM $(g-2)_\mu$?** [Stöckinger, Kaoru's talk]
- link to flavor physics and baryogenesis? [Standard Model fine??]
- mechanism for SUSY masses unknown [soft SUSY breaking mediated somehow?]

maximally blind mediation: mSUGRA [not a LHC paradigm!]

scalars: m_0 , fermions: $m_{1/2}$, tri-scalar term: A_0

plus $\text{sign}(\mu)$ and $\tan\beta$ in Higgs sector

- alternatives: gauge, anomaly, gaugino mediation ... ?
- ⇒ **measure spectrum at LHC instead**

LHC phenomenology: MSSM


- conjugate Higgs field not allowed
 - give mass to t and b ?
 - two Higgs doublets
- SUSY Higgs alone interesting
- ⇒ would be another talk...
- ⇒ **SUSY partners at LHC**

		spin	d.o.f.	
fermion	\hat{f}_L, \hat{f}_R	1/2	1+1	
→ sfermion	\hat{f}_L, \hat{f}_R	0	1+1	
gluon	\hat{G}_μ	1	n-2	
→ gluino	\hat{g}	1/2	2	Majorana
gauge bosons	γ, Z	1	2+3	
Higgs bosons	H^0, H^\pm, A^0	0	3	
→ neutralinos	$\hat{\chi}_j^0$	1/2	4 · 2	LSP?
gauge bosons	W^\pm	1	2 · 3	
Higgs bosons	H^\pm	0	2	
→ charginos	$\hat{\chi}_j^\pm$	1/2	2 · 4	
graviton	\hat{G}	2	2	
→ gravitino	\hat{G}	3/2	2	tough

Supersymmetry at LHC: 1

Inclusive: squarks and gluinos at Tevatron


- squarks, gluinos strongly interacting
 $p\bar{p} \rightarrow \tilde{q}\tilde{q}^*, \tilde{q}\tilde{g}, \tilde{g}\tilde{g}$ [best if $m(\tilde{q}) \sim m(\tilde{g})$]
 - large rates at hadron colliders
 - decays to jets and LSP
 $\tilde{g} \rightarrow \tilde{q}\bar{q}, \tilde{q}_L \rightarrow q\tilde{\chi}_2^0, \tilde{q}_R \rightarrow q\tilde{\chi}_1^0$
[additional jets and leptons possible]
 - gaugino mass unification assumed for details
- ⇒ we know inclusive jets plus LSP



Supersymmetry at LHC: 1


Inclusive: squarks and gluinos at Tevatron

- squarks, gluinos strongly interacting
 $p\bar{p} \rightarrow \tilde{q}\tilde{q}^*, \tilde{q}\tilde{g}, \tilde{g}\tilde{g}$ [best if $m(\tilde{q}) \sim m(\tilde{g})$]
 - large rates at hadron colliders
 - decays to jets and LSP
 $\tilde{g} \rightarrow \tilde{q}\bar{q}, \tilde{q}_L \rightarrow q\tilde{\chi}_2^0, \tilde{q}_R \rightarrow q\tilde{\chi}_1^0$
[additional jets and leptons possible]
 - gaugino mass unification assumed for details
- ⇒ **we know inclusive jets plus LSP**



When do we see SUSY-QCD?


- gluinos: strongly interacting Majorana fermions
 - first jet in gluino decay: q or \bar{q}
 - final-state leptons with both charges
- ⇒ **like-sign dileptons from $\tilde{g}\tilde{g}$** [Barger,...; Barnett,...; Baer,...]



Supersymmetry at LHC: 2

New physics at the LHC

- (1) possible discovery — signals for new physics, exclusion of parameter space
- (2) measurements
- (3) parameter studies




Supersymmetry at LHC: 2

New physics at the LHC

- (1) **possible discovery** — signals for new physics, exclusion of parameter space
 - (2) **measurements** — masses, cross sections, decays
 - (3) **parameter studies** — MSSM Lagrangean, SUSY breaking
- ⇒ approach independent of new physics model

Some SUSY signals at LHC [NLO: Prospino2]

- jets and \cancel{E}_T : $pp \rightarrow \tilde{q}\tilde{q}^*, \tilde{g}\tilde{g}, \tilde{q}\tilde{g}$
 - like-sign dileptons: $pp \rightarrow \tilde{g}\tilde{g}$
 - funny tops: $pp \rightarrow \tilde{t}_1\tilde{t}_1^*$
 - tri-leptons: $pp \rightarrow \tilde{\chi}_2^0\tilde{\chi}_1^-$
 $[\tilde{\chi}_2^0 \rightarrow \tilde{\ell}\tilde{\ell} \rightarrow \tilde{\chi}_1^0\ell\bar{\ell}; \tilde{\chi}_1^- \rightarrow \tilde{\chi}_1^0\ell\bar{\nu}]$
- ⇒ inclusive: similar to Tevatron
- ⇒ **exclusive: enough events for studies**





New physics mass measurements: 1

Spectra from cascade decays

- decay $\tilde{g} \rightarrow \tilde{b}\tilde{b} \rightarrow \tilde{\chi}_2^0 b\tilde{b} \rightarrow \mu^+ \mu^- b\tilde{b}\tilde{\chi}_1^0$ [better not via Z or to τ]
- cross sections some 100 pb [more than 3×10^7 events]
- thresholds & edges [$m_{\tilde{\ell}\ell}^2 < (m_{\tilde{\chi}_2^0}^2 - m_{\tilde{\ell}}^2)(m_{\tilde{\ell}}^2 - m_{\tilde{\chi}_1^0}^2)/m_{\tilde{\ell}}^2$]
- detector resolution, calibration, systematic errors, shape analysis, cross sections as input?

⇒ spectrum information from decay kinematics


[Hinchliffe,...;Allanach,...; not only SUSY: Meade & Reece]



New physics mass measurements: 1

Spectra from cascade decays


- decay $\tilde{g} \rightarrow \tilde{b}\tilde{b} \rightarrow \tilde{\chi}_2^0 b\tilde{b} \rightarrow \mu^+ \mu^- b\tilde{b}\tilde{\chi}_1^0$ [better not via Z or to τ]
- cross sections some 100 pb [more than 3×10^7 events]
- thresholds & edges [$m_{\tilde{\ell}\ell}^2 < (m_{\tilde{\chi}_2^0}^2 - m_{\tilde{\ell}}^2)(m_{\tilde{\ell}}^2 - m_{\tilde{\chi}_1^0}^2)/m_{\tilde{\ell}}^2$]
- detector resolution, calibration, systematic errors, shape analysis, cross sections as input?



⇒ **spectrum information from decay kinematics** [mass differences with smaller errors]

Glino mass from kinematic endpoints


- all decay jets b -tagged [Gjelsten, Miller, Osland]
 - most of time: cascade assumption correct
- ⇒ gluino mass to $\sim 1\%$
[theoretically defined?]



Squarks and gluinos always with many jets [TP, Rainwater, Skands]

- cascade studies sensitive to jet simulation?
 - matrix element $\tilde{g}\tilde{g}+2j$ and $\tilde{u}_L\tilde{g}+2j$ [$p_{T,j} > 100$ GeV]
 - compared with Pythia shower [tuned at Tevatron]
- ⇒ **QCD no killer for decay analyses** [the heavier the better]

σ [pb]	tt_{600}	$\tilde{g}\tilde{g}$	$\tilde{u}_L\tilde{g}$
σ_{0j}	1.30	4.83	5.65
σ_{1j}	0.73	2.89	2.74
σ_{2j}	0.26	1.09	0.85



New physics mass measurements: 2

Squarks and gluinos always with many jets [TP, Rainwater, Skands]

- cascade studies sensitive to jet simulation?
 - matrix element $\tilde{g}\tilde{g}+2j$ and $\tilde{u}_L\tilde{g}+2j$ [$p_{T,j} > 100$ GeV]
 - compared with Pythia shower [tuned at Tevatron]
- ⇒ **QCD no killer for decay analyses** [the heavier the better]

σ [pb]	$t\bar{t}_{600}$	$\tilde{g}\tilde{g}$	$\tilde{u}_L\tilde{g}$
σ_{0j}	1.30	4.83	5.65
σ_{1j}	0.73	2.89	2.74
σ_{2j}	0.26	1.09	0.85

SUSY plus jets: complex final states [Smadgraph: Cho, Hagiwara, Kanzaki, TP, Rainwater, Stelzer]


- Majoranas and fermion number violation in Madgraph
 - tested set of Feynman rules [400+ processes: Madgraph - Whizard (Reuter) - Sherpa (Schumann)]
 - implemented in Madevent [Alwall, Maltoni: Louvain group]
- ⇒ publicly available: <http://madgraph.phys.ucl.ac.be>

New physics spin measurements: 1

All new physics is hypothesis testing


- assume squark cascade observed
- ⇒ strongly interacting scalar?
- ⇒ straw-man model where squark is a fermion: universal extra dimensions

[Appelquist, Cheng, Dobrescu; Cheng, Matchev, Schmaltz; spectra degenerate —ignore; cross section larger —ignore]



Squark–slepton cascade [Smillie, Webber, Athanasiou, Lester]

- decay chain $\tilde{q} \rightarrow \tilde{\chi}_2^0 \rightarrow \tilde{\ell} \rightarrow \tilde{\chi}_1^0$
- trick 1: compare with KK q, Z, ℓ, γ
- trick 2: ‘invariant angles’
⇒ $\hat{m} = m_{j\ell} / m_{j\ell}^{\max}$ most promising [Barr]
- typically largest $pp \rightarrow \tilde{q}\tilde{g}$
- trick 3: production asymmetry $\tilde{q} : \tilde{q}^* \sim 2 : 1$
⇒ $\mathcal{A} = [\sigma(j\ell^+) - \sigma(j\ell^-)] / [\sigma(j\ell^+) + \sigma(j\ell^-)]$




Masses or spin or both? [Arkani-Hamed,...]

- masses from kinematic endpoints [use $m_{ej}, m_{\ell\ell}, m_{j\ell\ell} \dots$]
- spins from distributions between endpoints [endpoints identical in SUSY and UED]

New physics spin measurements: 2

Back to sign of SUSY-QCD

- like-sign dileptons indicate Majorana fermion?
 - always like-sign dileptons from bosonic gluon
- ⇒ show gluino fermionic
- ⇒ compare with usual straw man [UED-Madgraph: Alves]








Back to sign of SUSY-QCD

- like-sign dileptons indicate Majorana fermion?
- always like-sign dileptons from bosonic gluon
- ⇒ show gluino fermionic
- ⇒ compare with usual straw man [UED-Madgraph: Alves]

Gluino-bottom cascade [Alves, Eboli, TP]

- decay chain like for gluino mass
- compare with first KK g, q, Z, ℓ, γ
- replace initial-state asymmetry by b vs. \bar{b}
- independent of production channels
- asymmetry to write down:

$$A = [\sigma(bl^+) - \sigma(bl^-)] / [\sigma(bl^+) + \sigma(bl^-)]$$
- [still visible after cuts and smearing]





Back to sign of SUSY-QCD

- like–sign dileptons indicate Majorana fermion?
- always like–sign dileptons from bosonic gluon
- ⇒ show gluino fermionic
- ⇒ compare with usual straw man [UED–Madgraph: Alves]

Gluino–bottom cascade [Alves, Eboli, TP]

- exchange $\tilde{\ell}_{LR}$ in cascade
- test of lepton-ino couplings
- stau mixing [Choi, Hagiwara, Kim, Mawatari, Zerwas]
- purely hadronic ϕ_{bb} [TP, Plümacher, Reinartz]
- independent of weak decays
- sensitive to gluino/KK-gluon boost
- compare two SUSY hypotheses
- neutralino–sneutrino LSP [TP, Pradler, Steffen]

⇒ masses and spins accessible in decays at LHC



Theory output from LHC: SUSY parameters

- parameters: weak-scale Lagrangean [Sfitter: Lafaye, TP, Rauch, D Zerwas; Fittino; Harvard]
- measurements: masses or edges
branching fractions
cross sections
- errors: general correlation, statistics & systematics & theory
- problem in grid: huge phase space, local minimum?
problem in fit: domain walls, global minimum?

First go at problem

- ask a friend how SUSY is broken \Rightarrow mSUGRA
 - fit $m_0, m_{1/2}, A_0, \tan \beta, \text{sign}(\mu), y_t, \dots$
 - no problem, include indirect constraints [Ellis, Heinemeyer, Olive, Weiglein,...]
- \Rightarrow probability map as of today [Allanach, Lester, Weber]
- \Rightarrow best fit from LHC/ILC measurements

	SPS1a	Δ LHC masses	Δ LHC edges	Δ ILC	Δ LHC+ILC
m_0	100	3.9	1.2	0.09	0.08
$m_{1/2}$	250	1.7	1.0	0.13	0.11
$\tan \beta$	10	1.1	0.9	0.12	0.12
A_0	-100	33	20	4.8	4.3

Supersymmetric parameters: 1

Tilman Plehn

Supersymmetry

LHC Signals

Masses

Spins

Parameters

Statistics

Theory output from LHC: SUSY parameters

- parameters: weak-scale Lagrangean [Sfitter: Lafaye, TP, Rauch, D Zerwas; Fittino; Harvard]
- measurements: masses or edges
branching fractions
cross sections
- errors: general correlation, statistics & systematics & theory
- problem in grid: huge phase space, local minimum?
problem in fit: domain walls, global minimum?

MSSM instead of mSUGRA [TP, Lafaye, D Zerwas]

- (1) grid for closed subset
 - (2) fit of other parameters
 - (3) complete fit
 - LHC+ILC perfect [Weiglein etal]
- ⇒ too few measurements?
secondary minima? ...

	LHC	ILC	LHC+ILC	SPS1a
$\tan\beta$	10.22 ± 9.1	10.26 ± 0.3	10.06 ± 0.2	10
M_1	102.45 ± 5.3	102.32 ± 0.1	102.23 ± 0.1	102.2
M_2	578.67 ± 15	fix 500	588.05 ± 11	589.4
$M_{\tilde{T}L}$	fix 500	197.68 ± 1.2	199.25 ± 1.1	197.8
$M_{\tilde{T}R}$	129.03 ± 6.9	135.66 ± 0.3	133.35 ± 0.6	135.5
$M_{\tilde{b}L}$	198.7 ± 5.1	198.7 ± 0.5	198.7 ± 0.5	198.7
$M_{\tilde{g}L}$	498.3 ± 110	497.6 ± 4.4	521.9 ± 39	501.3
$M_{\tilde{t}R}$	fix 500	420 ± 2.1	411.73 ± 12	420.2
$M_{\tilde{b}R}$	522.26 ± 113	fix 500	504.35 ± 61	525.6
A_τ	fix 0	-202.4 ± 89.5	352.1 ± 171	-253.5
A_t	-507.8 ± 91	-501.95 ± 2.7	-505.24 ± 3.3	-504.9
A_b	-784.7 ± 35603	fix 0	-977 ± 12467	-799.4

Supersymmetric parameters: 2

Probability maps of new physics [Allanach, Roszkowski]

- Bayes' theorem: $p(m|d) = p(d|m) p(m)/p(d)$ [$p(d)$ through normalization]
 - Pythia/Herwig/Sherpa: data given a model $p(d|m) \sim |\mathcal{M}|^2$
 - theorist's prejudice: model $p(m)$
- ⇒ given measurements: (1) compute probability map $p(m|d)$ of parameter space
(2) rank local maxima

Weighted Markov chains [scanning algorithm for many dimensions: Rauch & TP]

- classical: produce representative set of spin states
compute average energy based on this reduced sample
- ⇒ map (chain) based on probability of a state
expensive energy function on sample
- BSM physics: produce map $p(m|d)$ of parameter points
evaluate same probability from (binned) density [Allanach,...; Baltz,...; Roszkowski,...]
- ⇒ phase-space MCMC approach: weighted chain [two bins with p_1 : 10 with 2 or 11 points]
- already for mSUGRA: MCMC resolution not sufficient
- ⇒ use additional probability maximization to rank maxima


Supersymmetric parameters: 2

Probability maps of new physics [Allanach, Roszkowski]

- Bayes' theorem: $p(m|d) = p(d|m) p(m)/p(d)$ $[p(d)$ through normalization]
 - Pythia/Herwig/Sherpa: data given a model $p(d|m) \sim |\mathcal{M}|^2$
 - theorist's prejudice: model $p(m)$
- ⇒ given measurements: (1) compute probability map $p(m|d)$ of parameter space
(2) rank local maxima

Toy model [Rauch & TP]

- test function $V(\vec{x})$ in 5 dimensions [general high-dimensional extraction tool]
- Sfitter output #1: probability map
Sfitter output #2: list of local maxima [best fit]



$V=74.929$ @ (655.00,253.72,347.83,348.57,349.59)

$V=59.972$ @ (850.04,224.99,650.00,649.99,654.56)

$V=58.219$ @ (849.97,225.01,587.08,650.01,650.02)

$V=25.110$ @ (750.00,749.99,450.00,450.01,450.01)

$V=16.042$ @ (245.45,253.44,552.51,542.58,544.75)

$V=12.116$ @ (350.70,650.40,650.36,650.40,650.38)

...



Supersymmetric parameters: 2

Probability maps of new physics [Allanach, Roszkowski]

- Bayes' theorem: $p(m|d) = p(d|m) p(m)/p(d)$ [$p(d)$ through normalization]
 - Pythia/Herwig/Sherpa: data given a model $p(d|m) \sim |\mathcal{M}|^2$
 - theorist's prejudice: model $p(m)$
- ⇒ given measurements: (1) compute probability map $p(m|d)$ of parameter space
(2) rank local maxima

mSUGRA with LHC measurements alone

- SPS1a kinematic edges with free m_t
- as of yesterday: Sfitter probability maps [Lafaye, TP, Rauch, D Zerwas]




Supersymmetric parameters: 2

Probability maps of new physics [Allanach, Roszkowski]

- Bayes' theorem: $p(m|d) = p(d|m) p(m)/p(d)$ [$p(d)$ through normalization]
 - Pythia/Herwig/Sherpa: data given a model $p(d|m) \sim |\mathcal{M}|^2$
 - theorist's prejudice: model $p(m)$
- ⇒ given measurements: (1) compute probability map $p(m|d)$ of parameter space
 (2) rank local maxima

MSSM with LHC measurements alone

- SPS1a kinematic edges and free m_t
- as of yesterday: Sfitter probability map [Lafaye, TP, Rauch, D Zerwas]



Statistics and phenomenology: 1

Problems in the spin extraction

- unclear if spin information available [top partners: Meade & Reece]

⇒ **when can we distinguish two hypotheses?**

Statistics: Neyman–Pearson lemma

- correct hypothesis m_1 : SUSY cascade
wrong hypothesis m_2 : UED cascade

⇒ likelihood ratio $p(d|m_1)/p(d|m_2)$ best estimator

[lowest probability to mistake right for fluctuation of wrong (type-II error)]

- likelihood $p(d|m) \sim |\mathcal{M}|^2$

⇒ **compute maximum statistical significance** [Cranmer, TP]


Statistics and phenomenology: 1

Problems in the spin extraction

- unclear if spin information available [top partners: Meade & Reece]
- ⇒ **when can we distinguish two hypotheses?**

Statistics: Neyman–Pearson lemma

- correct hypothesis m_1 : Higgs signal in WBF $H \rightarrow \mu\mu$ [TP, Rainwater]
 - wrong hypothesis m_2 : background
 - ⇒ likelihood ratio $p(d|m_1)/p(d|m_2)$ best estimator
[lowest probability to mistake right for fluctuation of wrong (type-II error)]
 - likelihood $p(d|m) \sim |\mathcal{M}|^2$
 - ⇒ **compute maximum statistical significance** [Cranmer, TP]
- $$CL_b(q) = \int_q^\infty dq' \rho_b(q') \quad [5\sigma \text{ is } CL_b = 2.85 \cdot 10^{-7}]$$



'Matrix element method' [CDF, D0; McElrath]

- compute likelihood of top events estimating $|\mathcal{M}|^2$
- maximize likelihood $p(d|SM, m_t)$ as function of m_t ...

Maximum significance for LHC signals

- example: combined n -event Poisson $[p(n|s+b) = e^{-(s+b)} (s+b)^n / n!]$

$$q = \log \frac{p(n|s+b)}{p(n|b)} = -s + n \log \left(1 + \frac{s}{b} \right) \longrightarrow - \sum_j s_j + \sum_j n_j \log \left(1 + \frac{s_j}{b_j} \right)$$

- phase space integration over $p(d|s, b) \sim |\mathcal{M}_{s,b}|^2$ [LEP-Higgs inspired]

$$q(\vec{r}) = -\sigma_s \mathcal{L} + \log \left(1 + \frac{|\mathcal{M}_s(\vec{r})|^2}{|\mathcal{M}_b(\vec{r})|^2} \right)$$

- probability distribution function via Fourier transform: $p_{s,b}(q)$
- \Rightarrow compute $CL_b(q)$

Maximum significance for LHC signals

- example: combined n -event Poisson $[p(n|s+b) = e^{-(s+b)} (s+b)^n / n!]$

$$q = \log \frac{p(n|s+b)}{p(n|b)} = -s + n \log \left(1 + \frac{s}{b} \right) \longrightarrow - \sum_j s_j + \sum_j n_j \log \left(1 + \frac{s_j}{b_j} \right)$$

- phase space integration over $p(d|s, b) \sim |\mathcal{M}_{s,b}|^2$ [LEP-Higgs inspired]

$$q(\vec{r}) = -\sigma_s \mathcal{L} + \log \left(1 + \frac{|\mathcal{M}_s(\vec{r})|^2}{|\mathcal{M}_b(\vec{r})|^2} \right)$$

- probability distribution function via Fourier transform: $\rho_{s,b}(q)$
- \Rightarrow compute $CL_b(q)$

Semi-realistic results [Cranmer & TP]

- irreducible & unsmeared

$$\sigma_{tot} = \int d\vec{r} M(\vec{r}) d\sigma(\vec{r})$$

- smearing $\Delta m_{\mu\mu}^{\text{width}} \ll \Delta m_{\mu\mu}^{\text{meas}}$ [unobserved dimensions]

$$\sigma_{tot} = \int d\vec{r}_\perp dr_m^* \int_{-\infty}^{\infty} dr_m M(\vec{r}) d\sigma(\vec{r}) W(r_m, r_m^*)$$

- acceptance cuts to reduce phase space... [bad measurements]
- \Rightarrow WBF $H \rightarrow \mu\mu$: 3.5σ in 300 fb^{-1}

Maximum significance for LHC signals

- example: combined n -event Poisson [$p(n|s+b) = e^{-(s+b)} (s+b)^n / n!$]

$$q = \log \frac{p(n|s+b)}{p(n|b)} = -s + n \log \left(1 + \frac{s}{b} \right) \longrightarrow - \sum_j s_j + \sum_j n_j \log \left(1 + \frac{s_j}{b_j} \right)$$

- phase space integration over $p(d|s, b) \sim |\mathcal{M}_{s,b}|^2$ [LEP-Higgs inspired]

$$q(\vec{r}) = -\sigma_s \mathcal{L} + \log \left(1 + \frac{|\mathcal{M}_s(\vec{r})|^2}{|\mathcal{M}_b(\vec{r})|^2} \right)$$

- probability distribution function via Fourier transform: $\rho_{s,b}(q)$
- \Rightarrow compute $CL_b(q)$

Estimate power of analyses [Cranmer, TP, Reuter]

- numerical evaluation instantaneous [sanity-check neural nets?]
- implementation into Whizard on the way
- cannot claim if search is possible
- **but can tell if search is impossible**

New physics at the LHC

A lot has been done

- higher-order calculations
- improved background estimates
- web-based signal event generators
- mass and spin measurements
- parameter extraction/probability maps
-

A lot is still left to do

- more higher-order calculations
 - even better background estimates
 - QCD effects on new physics measurements
 - scans of high-dimensional parameter spaces
 - statistics tools for phenomenologists
 -
- ⇒ even exciting times require serious work

**New Methods
for New Physics**

Tilman Plehn

Supersymmetry

LHC Signals

Masses

Spins

Parameters

Statistics