

UV Completions of the Standard Model at LHC

Tilman Plehn

University of Edinburgh

Saclay, 3/2008

Outline

TeV scale in the LHC era

Decays and masses

Decays and underlying parameters

Spins and jets I

Spins and jets II

Predictive extra dimensions

Standard–Model effective theory

Remember the Standard Model

- gauge theory $SU(3) \times SU(2) \times U(1)$
 - massless $SU(3)$ and $U(1)$ gauge bosons
massive $SU(2)$ gauge bosons [spontaneous symmetry breaking]
 - massive Dirac fermions [via Yukawas]
 - perturbatively renormalizable Lagrangian [no 1/M terms]
 - one missing piece: Higgs [fundamental? minimal? mass?]
- ⇒ truly fundamental theory

How complete experimentally?


- dark matter? [solid evidence! — for weak–scale new physics?]
 - $(g - 2)_\mu$? [possible evidence for weak–scale new physics, review by Dominik]
 - quark mixing — flavor physics? [new operators above 10^4 GeV?]
 - neutrino masses and mixing? [see-saw at 10^{11} GeV?]
 - matter–antimatter asymmetry? [universe mostly matter]
 - gauge–coupling unification? [almost perfect, but proton stable]
 - gravity? [mostly negligible but perturbatively non-renormalizable]
- ⇒ cut-off scale unavoidable: SM effective theory

Standard–Model effective theory

Remember the Standard Model

- gauge theory $SU(3) \times SU(2) \times U(1)$
- massless $SU(3)$ and $U(1)$ gauge bosons
massive $SU(2)$ gauge bosons [spontaneous symmetry breaking]
- massive Dirac fermions [via Yukawas]
- perturbatively renormalizable Lagrangian [no $1/M$ terms]
- one missing piece: Higgs [fundamental? minimal? mass?]

⇒ truly fundamental theory



How consistent theoretically?

- problem of light Higgs: mass driven to cutoff of effective Standard Model
$$\delta m_H^2 \propto g^2(2m_W^2 + m_Z^2 + m_t^2 - 4m_t^2) \Lambda^2$$
- cancelled by finely tuned counter term? [ugly, against spirit of symmetries]
- why fundamental Higgs at all, if such poor high-energy behavior?
- tied in with new physics at TeV scale:
 - supersymmetry [my favorite]
 - extra dimensions [cool idea]
 - little Higgs [old idea, now working]
 - no fundamental Higgs [hard with top mass]

⇒ TeV scale: beautiful ideas — complicated realistic models

Effective Standard Model in the LHC era

UV completions from the LHC [Uli Baur's rule: 'there is always new physics at higher scales']

- find light Higgs?
- find new physics stabilizing Higgs mass?
- see dark-matter candidate?

Particle theory in the LHC era

- model-independent analyses not helpful to understand TeV scale
 - early data: probably with little information [just a little pheno game]
 - understood data: testable TeV-scale models [e.g. Higgs sector vs. underlying theory?]
 - link to other observations [DM+Tevatron: Hooper, TP, Valinotto]
 - reconstruction of Lagrangian
- ⇒ **LHC theory without believing any model**

Effective Standard Model in the LHC era

UV completions from the LHC [Uli Baur's rule: 'there is always new physics at higher scales']

- find light Higgs?
- find new physics stabilizing Higgs mass?
- see dark-matter candidate?

Particle theory in the LHC era


- model-independent analyses not helpful to understand TeV scale
- early data: probably with little information [just a little pheno game]
- understood data: testable TeV-scale models [e.g. Higgs sector vs. underlying theory?]
- link to other observations [DM+Tevatron: Hooper, TP, Valinotto]
- reconstruction of Lagrangian

⇒ LHC theory without believing any model


Role of LHC

- beyond inclusive searches [that was Tevatron]
millions of new strongly interacting particles

⇒ (1) aim at underlying theory
(2) try to survive QCD [skipping today]



TeV-scale supersymmetry



Supersymmetry

- give each Standard-Model particle a partner [with different spin, including strong interactions]
 - symmetry obviously broken by masses [soft breaking, mechanism unknown]
 - assume dark matter, stable lightest partner [R parity]
- ⇒ **measure BSM spectrum with missing energy at LHC**

LHC searches: MSSM


- SUSY-Higgs alone interesting...
...but not conclusive
...and another talk
- ⇒ **list of SUSY partners**

		spin	d.o.f.	
fermion	f_L, f_R	1/2	1+1	
→ sfermion	\tilde{f}_L, \tilde{f}_R	0	1+1	
gluon	G_μ	1	n-2	
→ gluino	\tilde{g}	1/2	2	Majorana
gauge bosons	γ, Z	1	2+3	
Higgs bosons	h^0, H^0, A^0	0	3	
→ neutralinos	$\tilde{\chi}_i^0$	1/2	4 · 2	LSP
gauge bosons	W^\pm	1	2 · 3	
Higgs bosons	H^\pm	0	2	
→ charginos	$\tilde{\chi}_i^\pm$	1/2	2 · 4	

Decays and masses





Cascade decays [Atlas-TDR, Cambridge]

- if new particles strongly interacting and LSP weakly interacting
- like Tevatron: jets + missing energy [rates tough because of QCD]
- cascade kinematics [$10^7 \dots 10^8$ events]



$$0 < m_{\mu\mu}^2 < \frac{m_{\tilde{\chi}_2^0}^2 - m_{\tilde{\ell}}^2}{m_{\tilde{\ell}}} \frac{m_{\tilde{\ell}}^2 - m_{\tilde{\chi}_1^0}^2}{m_{\tilde{\ell}}}$$


⇒ new-physics mass spectrum from cascade kinematics



Decays and masses

Cascade decays [Atlas-TDR, Cambridge]

- if new particles strongly interacting and LSP weakly interacting
- like Tevatron: jets + missing energy [rates tough because of QCD]
- cascade kinematics [$10^7 \dots 10^8$ events]




$$0 < m_{\mu\mu}^2 < \frac{m_{\tilde{\chi}_2^0}^2 - m_{\tilde{\ell}}^2}{m_{\tilde{\ell}}} \frac{m_{\tilde{\ell}}^2 - m_{\tilde{\chi}_1^0}^2}{m_{\tilde{\ell}}}$$

⇒ **new-physics mass spectrum from cascade kinematics**

Glucino decay [Gjelsten, Miller, Osland, Raklev]

- no problem: additional jets [Rainwater, TP, Skands; Alwall, Wacker,...]
- no problem: off-shell effects [Catpiss collaboration; Kauer & Rainwater]
- all decay jets *b* quarks [otherwise dead by QCD]
- limited by jet energy scale
- alternative methods?

⇒ **why physical masses?**



Decays and underlying parameters

From kinematics to weak-scale parameters [Fittino; SFitter: Lafaye, TP, Rauch, Zerwas]

- parameters: weak-scale Lagrangian
- measurements: kinematic endpoints, branching ratios, rates [Prospino2]
 B decays, $(g - 2)_\mu$, dark matter, e-w precision data...
- errors: general correlation, statistics & systematics & theory [flat theory errors!]
- problem in grid/fit: no local/global maximum
problem in physics: secondary maxima

Probability maps of new physics [Baltz,...; Roszkowski,...; Allanach,...; SFitter]

- fully exclusive likelihood map $p(d|m)$ over m [hard part]
- LHC problem: remove directions
- Bayesian: $p(m|d) \sim p(d|m) p(m)$ with theorists' bias $p(m)$ [cosmology, BSM]
frequentist: best-fitting point $\max_m p(d|m)$ [flavor]
- LHC era: (1) compute high-dimensional map $p(d|m)$
(2) find and rank local maxima in $p(d|m)$
(3) Bayesian-frequentist dance to reduce dimensions

Decays and underlying parameters

From kinematics to weak-scale parameters [Fittino; SFitter: Lafaye, TP, Rauch, Zerwas]

- parameters: weak-scale Lagrangian
- measurements: kinematic endpoints, branching ratios, rates [Prospino2]
 B decays, $(g - 2)_\mu$, dark matter, e-w precision data...
- errors: general correlation, statistics & systematics & theory [flat theory errors!]
- problem in grid/fit: no local/global maximum
problem in physics: secondary maxima


Toy model: MSUGRA map from LHC [LHC endpoints with free y_t]

- weighted Markov chains: several times faster [similar to: Ferrenberg & Swendsen]

$$P_{\text{bin}}(p \neq 0) = \frac{N}{\sum_{i=1}^N 1/p}$$

- SFitter output #1: fully exclusive likelihood map
- SFitter output #2: ranked list of local maxima

⇒ **strong correlations even in MSUGRA**





Underlying parameters

MSSM map at LHC [for details ask Dirk]

- shifting from 6D to 19D parameter space [killing grids, Minuit, laptop-style fits...]
- SFitter outputs still the same, but best points degenerate
- e.g. 3 neutralinos observed

Bayesian pdf noisy
profile likelihood no pdf





⇒ no golden approach to BSM statistics

Underlying parameters

MSSM map at LHC [for details ask Dirk]

- shifting from 6D to 19D parameter space [killing grids, Minuit, laptop-style fits...]
- SFitter outputs still the same, but best points degenerate
- e.g. 3 neutralinos observed


Bayesian pdf noisy
profile likelihood no pdf



⇒ no golden approach to BSM statistics

MSSM map in LHC era [SFitter+friends]

- LHC: $\text{sign}(\mu)$ unknown and $\tan\beta$ extraction believe-based





Underlying parameters

MSSM map at LHC [for details ask Dirk]

- shifting from 6D to 19D parameter space [killing grids, Minuit, laptop-style fits...]
- SFitter outputs still the same, but best points degenerate
- e.g. 3 neutralinos observed

Bayesian pdf noisy
profile likelihood no pdf



⇒ no golden approach to BSM statistics

MSSM map in LHC era [Sfitter+friends]

- LHC: $\text{sign}(\mu)$ unknown and $\tan \beta$ extraction believe-based
- $\tan \beta$ and $\text{sign}(\mu)$ from $(g-2)_\mu$ or $B_s \rightarrow \mu\mu$ [Les Houches 2007: Alexander et al]
- $(g-2)_\mu$ very promising
- B_s decays killed by non-pert. QCD

	LHC	LHC $\otimes (g-2)$	SPS1a
$\tan \beta$	10.0 ± 4.5	10.3 ± 2.0	10.0
M_1	102.1 ± 7.8	102.7 ± 5.9	103.1
M_2	193.3 ± 7.8	193.2 ± 5.8	192.9
M_3	577.2 ± 14.5	578.2 ± 12.1	577.9
$M_{\tilde{\mu}L}$	193.2 ± 8.8	194.0 ± 6.8	194.4
$M_{\tilde{\mu}R}$	135.0 ± 8.3	135.6 ± 6.3	135.8
$M_{\tilde{q}L}$	524.6 ± 14.5	525.5 ± 10.6	526.6
$M_{\tilde{q}R}$	507.3 ± 17.5	507.6 ± 15.8	508.1
μ	350.5 ± 14.5	352.5 ± 10.8	353.7

Underlying parameters

MSSM map at LHC [for details ask Dirk]


- shifting from 6D to 19D parameter space [killing grids, Minuit, laptop-style fits...]
 - SFitter outputs still the same, but best points degenerate
- ⇒ **no golden approach to BSM statistics**

MSSM map in LHC era [SFitter+friends]

- LHC: $\text{sign}(\mu)$ unknown and $\tan\beta$ extraction believe-based
- $\tan\beta$ and $\text{sign}(\mu)$ from $(g-2)_\mu$ or $B_s \rightarrow \mu\mu$ [Les Houches 2007: Alexander et al]
- $(g-2)_\mu$ very promising
- B_s decays killed by non-pert. QCD

Fundamental theory [SFitter + Kneur]



- SUSY breaking?
 - unification, GUT?
 - scale-invariant sum rules? [Cohen, Schmalz]
 - new & crucial: renormalization group bottom-up
- ⇒ **LHC sensitive to UV models**



Spins from cascades

Supersymmetry at Tevatron/LHC: Majorana gluinos [Barger,...; Barnett,...; Baer,...]

- LHC: first jet (q or \bar{q}) fixes lepton charge
 - same-sign dileptons in 1/2 of events
 - similar: t -channel gluino in $pp \rightarrow \tilde{q}\tilde{q}$
- \Rightarrow **gluino = like-sign dileptons in SUSY sample**




Spins from cascades

Supersymmetry at Tevatron/LHC: Majorana gluinos [Barger,...; Barnett,...; Baer,...]

- LHC: first jet (q or \bar{q}) fixes lepton charge
 - same–sign dileptons in 1/2 of events
 - similar: t -channel gluino in $pp \rightarrow \tilde{q}\tilde{q}$
- ⇒ **gluino = like–sign dileptons in SUSY sample**

Loop hole: gluino is Majorana if fermion [Alves, Eboli, TP]

- start with mass–measurement cascade
- now: physics between the endpoints
- model–independent analysis difficult [Barr, Lester, Smillie, Webber]



Spins from cascades


Supersymmetry at Tevatron/LHC: Majorana gluinos [Barger,...; Barnett,...; Baer,...]

- LHC: first jet (q or \bar{q}) fixes lepton charge
 - same-sign dileptons in 1/2 of events
 - similar: t -channel gluino in $pp \rightarrow \tilde{q}\tilde{q}$
- ⇒ **gluino = like-sign dileptons in SUSY sample**

Loop hole: gluino is Majorana if fermion [Alves, Eboli, TP]

- start with mass-measurement cascade
- now: physics between the endpoints
- model-independent analysis difficult [Barr, Lester, Smillie, Webber]

- UV-completion hypotheses tests instead
 - 'gluino' a boson: universal extraD
 - compare SUSY vs. KK g, b, Z, l, γ
 - simple distributions $\Delta\phi_{bb}$ [3-body decays: Csaki,...]
- ⇒ **gluino = fermion with like-sign dileptons**



Spins from cascades


Supersymmetry at Tevatron/LHC: Majorana gluinos [Barger,...; Barnett,...; Baer,...]

- LHC: first jet (q or \bar{q}) fixes lepton charge
 - same-sign dileptons in 1/2 of events
 - similar: t -channel gluino in $pp \rightarrow \tilde{q}\tilde{q}$
- ⇒ **gluino = like-sign dileptons in SUSY sample**

Loop hole: gluino is Majorana if fermion [Alves, Eboli, TP]

- start with mass-measurement cascade
- now: physics between the endpoints
- model-independent analysis difficult [Barr, Lester, Smillie, Webber]


- UV-completion hypotheses tests instead
 - 'gluino' a boson: universal extraD
 - compare SUSY vs. KK g, b, Z, l, γ
 - simple distributions $\Delta\phi_{bb}$ [3-body decays: Csaki,...]
- ⇒ **gluino = fermion with like-sign dileptons**
- sensitive to model's details
- ⇒ **LHC requiring testable TeV-scale models**



Spins from forward jets

Beyond cascades: spin of LSP [Alwall, TP, Rainwater]


- hypotheses: like–sign charginos (Majorana neutralino)
like–sign scalars (stable scalars)
like–sign vector bosons (little–Higgs inspired)
- chargino decay/kinematics not used, jet correlations the key



Spins from forward jets

Beyond cascades: spin of LSP [Allwall, TP, Rainwater]

- hypotheses: like–sign charginos (Majorana neutralino)
 like–sign scalars (stable scalars)
 like–sign vector bosons (little–Higgs inspired)
- chargino decay/kinematics not used, jet correlations the key




Like-sign scalars instead of fermions

- charged Higgs in 2HDM [neutral Higgs: TP, Rainwater, Zeppenfeld; Buszello, Marquard, v.d.Bij]
- W radiated off quarks [Goldstone coupling to Higgs]

$$P_T(x, p_T) \sim \frac{1 + (1-x)^2}{2x} \frac{1}{p_T^2}$$

$$P_L(x, p_T) \sim \frac{(1-x)^2}{x} \frac{m_W^2}{p_T^4}$$


⇒ scalars with softer $p_{T,j}$



Spins from forward jets

Beyond cascades: spin of LSP [Alwall, TP, Rainwater]

- hypotheses: like-sign charginos (Majorana neutralino)
 like-sign scalars (stable scalars)
 like-sign vector bosons (little-Higgs inspired)
- chargino decay/kinematics not used, jet correlations the key



Like-sign scalars instead of fermions

- charged Higgs in 2HDM [neutral Higgs: TP, Rainwater, Zeppenfeld; Buszello, Marquard, v.d.Bij]
- W radiated off quarks [Goldstone coupling to Higgs]

$$P_T(x, p_T) \sim \frac{1 + (1-x)^2}{2x} \frac{1}{p_T^2}$$


$$P_L(x, p_T) \sim \frac{(1-x)^2}{x} \frac{m_W^2}{p_T^4}$$

⇒ scalars with softer $p_{T,j}$

Like-sign vectors instead of fermions

- little-Higgs inspired [T -type parity]
- start with copy of SM, heavy W', Z', H', f'
 [H' necessary for unitarity, but irrelevant at LHC]
- Lorentz structure in angle between jets


⇒ vectors with peaked $\Delta\phi_{jj}$



Spins from forward jets

Beyond cascades: spin of LSP [Alwall, TP, Rainwater]

- hypotheses: like-sign charginos (Majorana neutralino)
like-sign scalars (stable scalars)
like-sign vector bosons (little-Higgs inspired)
- chargino decay/kinematics not used, jet correlations the key



Like-sign scalars instead of fermions

- charged Higgs in 2HDM [neutral Higgs: TP, Rainwater, Zeppenfeld; Buszello, Marquard, v.d.Bij]
- W radiated off quarks [Goldstone coupling to Higgs]

$$P_T(x, p_T) \sim \frac{1 + (1-x)^2}{2x} \frac{1}{p_T^2}$$


$$P_L(x, p_T) \sim \frac{(1-x)^2}{x} \frac{m_W^2}{p_T^4}$$

⇒ scalars with softer $p_{T,j}$

Heavy fermions in little-Higgs models

- part of unitary UV completion [Englert, Zeppenfeld]
- huge effects on distributions [strongly interacting W s]

⇒ LHC requiring testable TeV-scale models



Predictive extra dimensions

Elegant solution to hierarchy problem [Antoniadis, Arkani-Hamed, Dimopoulos, Dvali]

- highest scale: Planck scale $G_N \sim 1/M_{\text{Planck}}^2$ [$M_{\text{Planck}} \sim 10^{19}$ GeV]
- Einstein–Hilbert action in $4 + n$ dimensions [on torus — periodic boundaries]

$$\int d^4x \sqrt{|g|} M_{\text{Planck}}^2 R \rightarrow \int d^{4+n}x \sqrt{|g|} M_D^{2+n} R = (2\pi r)^n \int d^4x \sqrt{|g|} M_D^{2+n} R$$

$$M_{\text{Planck}} = M_D (2\pi r M_D)^{n/2} \gg M_D \sim 1 \text{ TeV}$$

- to get numbers right: $r = 10^{12}, 10^{-3}, \dots 10^{-11}$ m for $n = 1, 2, \dots 6$
- ⇒ **fundamental Planck scale at TeV**

Kaluza–Klein gravitons [Giudice, Ratazzi, Wells]

- periodic boundaries: Fourier–transform in extra dimensions [QCD massless]

$$(\square + m_k^2) G_{\mu\nu}^{(k)} = -\frac{T_{\mu\nu}}{M_{\text{Planck}}} \quad \delta m \sim \frac{1}{r} = 2\pi M_D \left(\frac{M_D}{M_{\text{Planck}}} \right)^{2/n} \lesssim 0.05 \text{ GeV}$$

- KK tower of single gravitons, each coupled as $1/M_{\text{Planck}}$
- IR spectrum: constraints from supernova cooling
- UV spectrum: LHC effects at TeV scale [Giudice, Strumia, TP; cosmic rays]

Predictive extra dimensions

Hope for collider searches [Giudice, Rattazzi, Wells; Han, Lykken, Zhang]

- real radiation of continuous KK tower

$$\sigma^{\text{tower}} \sim \sigma^{\text{graviton}} \int dm S_{n-1} m^{n-1} r^n = \sigma^{\text{graviton}} \int dm \frac{S_{n-1} m^{n-1}}{(2\pi M_D)^n} \left(\frac{M_{\text{Planck}}}{M_D} \right)^2$$

- higher-dimensional operator from virtual gravitons [UV dominated]

$$\mathcal{A}(s; m) = \frac{1}{M_{\text{Planck}}^2} T_{\mu\nu} T^{\mu\nu} \frac{1}{s - m^2} \rightarrow \frac{S_{n-1}}{2M_D^4} \left(\frac{\Lambda}{M_D} \right)^{n-2}$$

- KK tower coupling with $1/M_D$ at LHC
- LHC rates cut-off dependent

⇒ **KK effective theory poor at LHC, UV completion needed**



Predictive extra dimensions

Renormalization flow of gravity [Reuter; Litim; Wetterich;...]

- Einstein–Hilbert action with running $G(\mu)$ and $\Lambda(\mu)$
- dimensionless coupling $g(\mu) = G(\mu) \mu^{2+n} = G_0 Z_G^{-1}(\mu) \mu^{2+n}$
- attractive finite UV fixed point [anomalous dimension: $\eta = -\mu \partial_\mu \log Z_G \propto g$]

$$\mu \frac{\partial}{\partial \mu} g_*(\mu) = (2 + n + \eta(g_*)) g_*(\mu) = 0 \quad \text{for} \quad g_* \neq 0 \quad \eta(g_*) = -2 - n$$

- gravity asymptotically free in UV $G(\mu) \sim g_* \mu^{-(2+n)}$
- ⇒ **coupling weak enough for finite LHC predictions?**

Predictive extra dimensions

Renormalization flow of gravity [Reuter; Litim; Wetterich;...]

- Einstein–Hilbert action with running $G(\mu)$ and $\Lambda(\mu)$
- dimensionless coupling $g(\mu) = G(\mu) \mu^{2+n} = G_0 Z_G^{-1}(\mu) \mu^{2+n}$
- attractive finite UV fixed point [anomalous dimension: $\eta = -\mu \partial_\mu \log Z_G \propto g$]

$$\mu \frac{\partial}{\partial \mu} g_*(\mu) = (2 + n + \eta(g_*)) g_*(\mu) = 0 \quad \text{for} \quad g_* \neq 0 \quad \eta(g_*) = -2 - n$$

- gravity asymptotically free in UV $G(\mu) \sim g_* \mu^{-(2+n)}$
- ⇒ **coupling weak enough for finite LHC predictions?**

UV–safe gravity [Weinberg (1979)]

- gravity non–fundamental effective theory $G \propto 1/M_D^{2+n}?$
- 't Hooft's perturbative renormalizability: finite number of counter terms
- Wilson's renormalizability: no unphysical UV divergences
- consistent theory beyond perturbation theory [no ghosts: Weinberg & Gomez]
- fixed point likely to $\sqrt{|g|} R^{\frac{D}{2}}$ and including matter [no proof; not perturbative series]
- great idea for gravity — great for LHC

Predictive extra dimensions

UV-completed graviton production [Litim & TP]

– form factor for $G(\mu)$ [Hewett & Rizzo] $\frac{1}{M_D^{2+n}} \longrightarrow \frac{1}{M_D^{2+n}} \left[1 + \left(\frac{\sqrt{s}}{aM_D} \right)^{2+n} \right]^{-1}$

– alternative: changing anomalous dimension of graviton [QCD inspired]


$$P(s, m) = \frac{1}{s + m^2} \longrightarrow \frac{M_D^{n+2}}{(s + m^2)^{n/2+2}} \quad \text{around } m \sim M_D$$

– integration kernel after integration over m_{KK}

$$\frac{1}{M_D^{2+n}} \int_0^\infty \frac{dm}{m^{n-1}} P(s, m) = \frac{1}{n-2} \frac{1}{M_D^4} \left(\frac{aM_D}{M_D} \right)^{n-2} \left[1 + \frac{n-2}{4} \right] \left[1 + \mathcal{O} \left(\frac{s}{M_D^2} \right) \right]$$

– $\sqrt{s} > M_D$: kernel only function of \sqrt{s} [matching at M_D ?, black-hole solutions?]

⇒ UV-safe gravity safe at LHC



Predictive extra dimensions

UV-completed graviton production [Litim & TP]

– form factor for $G(\mu)$ [Hewett & Rizzo] $\frac{1}{M_D^{2+n}} \rightarrow \frac{1}{M_D^{2+n}} \left[1 + \left(\frac{\sqrt{s}}{aM_D} \right)^{2+n} \right]^{-1}$

– alternative: changing anomalous dimension of graviton [QCD inspired]

$$P(s, m) = \frac{1}{s + m^2} \rightarrow \frac{M_D^{n+2}}{(s + m^2)^{n/2+2}} \quad \text{around } m \sim M_D$$

– integration kernel after integration over m_{KK}

$$\frac{1}{M_D^{2+n}} \int_0^\infty \frac{dm}{m^{n-1}} P(s, m) = \frac{1}{n-2} \frac{1}{M_D^4} \left(\frac{aM_D}{M_D} \right)^{n-2} \left[1 + \frac{n-2}{4} \right] \left[1 + \mathcal{O} \left(\frac{s}{M_D^2} \right) \right]$$

– $\sqrt{s} > M_D$: kernel only function of \sqrt{s} [matching at M_D ?, black-hole solutions?]

⇒ **UV-safe gravity safe at LHC**

String theory as UV completion [e.g. Cullen, Perelstein, Peskin]

– Veneziano form factor

$$\frac{\Gamma(1 - \alpha' s) \Gamma(1 - \alpha' t)}{\Gamma(1 - \alpha' (s + t))} = \frac{\Gamma(1 - s/M_S^2) \Gamma(1 - t/M_S^2)}{\Gamma(1 - (s + t)/M_S^2)} = 1 - \frac{\pi^2}{6} \frac{st}{M_S^4} + \mathcal{O}(M_S^{-6})$$

– string resonances in UV: $\sqrt{n} M_S$

⇒ **quantum gravity testable at LHC**

New physics at the LHC

Physics in the LHC era

- understand e-w symmetry breaking
- confirm new physics [dark matter]
- complete Standard Model



New physics at the LHC

Physics in the LHC era

- understand e-w symmetry breaking
- confirm new physics [dark matter]
- complete Standard Model

LHC physics is fun physics!

- look for solid new–physics signals
 - measure weak–scale Lagrangian
 - determine fundamental physics

 - construct testable new–physics hypotheses
 - implement into realistic simulations
 - avoid getting killed by QCD
- ⇒ **LHC + testable TeV-scale models: more than discovery machine!**



**UV Completions of
the Standard
Model at LHC**

Tilman Plehn

TeV scale

Masses

Parameters

Spins & jets I

Spins & jets II

Extra dimensions